

Tobacco in Australia

Facts & Issues

Relevant news and research

12.4 Emissions from tobacco products

Last updated October 2024

Research:.....	3
12.4 Emissions from tobacco products.....	3
12.4.1 What is tobacco smoke?	4
12.4.1.1 Particles and gases in tobacco smoke.....	6
12.4.1.2 Mainstream smoke, sidestream smoke and secondhand smoke.....	7
12.4.2 What happens when tobacco products are burned?	10
12.4.2.1 Pyrolysis	10
12.4.2.2 Combustion.....	11
12.4.2.3 Formation of smoke in cigarettes	12
12.4.2.4 Formation of smoke in other types of combusted tobacco products	12
12.4.3 Chemicals that are emitted from combusted tobacco products.....	14
12.4.3.1 Nicotine and other alkaloids	21
12.4.3.2 Aldehydes.....	26
12.4.3.3 Aromatic amines	29
12.4.3.4 Hydrocarbons.....	30
12.4.3.5 Heterocyclic aromatic amines.....	31
12.4.3.6 Polycyclic aromatic hydrocarbons (PAH)	31
12.4.3.7 N-nitrosamines.....	34
12.4.3.8 Phenols.....	39
12.4.3.9 Other carbon-based (organic) compounds.....	40
12.4.3.10 Inorganics.....	41
12.4.3.11 Metals, metalloids and radionuclides.....	43
12.4.3.12 Additives and flavours (refers to section 12.6).....	49

12.4.4 Emission from tobacco that is not combusted (set alight)	49
12.4.4.1 Emissions from waterpipes	49
12.4.4.2 Emissions from heated tobacco ('heat-not-burn') products	51
12.4.4.3 Emission from smokeless tobacco	52
News:	53
12.4 Emissions from tobacco products.....	53
12.4.1 What is tobacco smoke?	54
12.4.1.1 Particles and gases in tobacco smoke.....	54
12.4.1.2 Mainstream smoke, sidestream smoke and secondhand smoke.....	54
12.4.2 What happens when tobacco products are burned?	54
12.4.2.1 Pyrolysis	54
12.4.2.2 Combustion.....	54
12.4.2.3 Formation of smoke in cigarettes	54
12.4.2.4 Formation of smoke in other types of combusted tobacco products	54
12.4.3 Chemicals that are emitted from combusted tobacco products.....	54
12.4.3.1 Nicotine and other alkaloids	55
12.4.3.2 Aldehydes.....	55
12.4.3.3 Aromatic amines	55
12.4.3.4 Hydrocarbons.....	55
12.4.3.5 Heterocyclic aromatic amines.....	55
12.4.3.6 Polycyclic aromatic hydrocarbons (PAH)	55
12.4.3.7 N-nitrosamines.....	55
12.4.3.8 Phenols.....	55
12.4.3.9 Other carbon-based (organic) compounds.....	55
12.4.3.10 Inorganics.....	56
12.4.3.11 Metals, metalloids and radionuclides.....	56
12.4.3.12 Additives and flavours (refers to section 12.6).....	56
12.4.4 Emission from tobacco that is not combusted (set alight)	56
12.4.4.1 Emissions from waterpipes	56
12.4.4.2 Emissions from heated tobacco ('heat-not-burn') products	56
12.4.4.3 Emission from smokeless tobacco	56

Research:

12.4 Emissions from tobacco products

Bos PMJ, Soeteman-Hernandez LG, and Talhout R. Risk assessment of components in tobacco smoke and e-cigarette aerosols: a pragmatic choice of dose metrics. *Inhalation Toxicology*, 2021; 33(3):81-95. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33876709>.

Pang YQ, Chen XJ, Li XY, Zhang HF, Jiang XY, et al. Magnetic solid phase extraction of tobacco-specific N-nitrosamines using magnetic graphene composite as sorbent. *Journal of Separation Science*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31347241>.

Ozdemir E, Es H, Demir M, and Uzun I. Forensic medical evaluation of deaths resulting from inhalation of cigarette lighter refill fuel in Turkey. *Legal Medicine (Tokyo, Japan)*, 2017; 24:1-6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28081784>.

Morabito JA, Holman MR, Ding YS, Yan X, Chan M, et al. The use of charcoal in modified cigarette filters for mainstream smoke carbonyl reduction. *Regulatory Toxicology and Pharmacology*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28238852>.

Baumung C, Rehm J, Franke H, and Lachenmeier DW. Comparative risk assessment of tobacco smoke constituents using the margin of exposure approach: the neglected contribution of nicotine. *Sci Rep*, 2016; 6:35577. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27759090>.

Administration FaD. "Harmful and Potentially Harmful Constituents" in Tobacco Products as Used in Section 904(e) of the Federal Food, Drug, and Cosmetic Act (Revised)*. U.S. Department of Health and Human Services, Food and Drug Administration, Center for Tobacco Products, 2016. Available from: <http://www.fda.gov/TobaccoProducts/Labeling/ProductsIngredientsComponents/ucm20035927.htm>.

Belushkin M, Jaccard G, and Kondylis A. Considerations for comparative tobacco product assessments based on smoke constituent yields. *Regulatory Toxicology and Pharmacology*, 2015; 73(1):105–13. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26140819>.

World Health Organization. The scientific basis of tobacco product regulation: Second report of a WHO study group. Geneva: WHO, 2008. Available from: https://apps.who.int/iris/handle/10665/43997?search-result=true&query=scientific+basis+for+tobacco+product+regulation&scope=&rpp=10&sort_by=score&order=desc.

World Health Organization. The scientific basis for product regulation. Technical Report Series, 945. Geneva: World Health Organisation, 2007. Available from: http://www.who.int/tobacco/global_interaction/tobreg/who_tsr.pdf.

Giess O and Kotzias D. Tobacco, Cigarettes and Cigarette Smoke: An Overview. Luxembourg: Institute for Health and Consumer Protection, European Commission, Joint Research Centre., 2007. Available from: <https://publications.jrc.ec.europa.eu/repository/handle/JRC37472>.

Ding YS, Yan XJ, Jain RB, Lopp E, Tavakoli A, et al. Determination of 14 polycyclic aromatic hydrocarbons in mainstream smoke from U.S. brand and non-U.S. brand cigarettes. *Environmental*

Science & Technology, 2006; 40(4):1133-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/16572766>

Adams JD, O'Mara-Adams KJ, and Hoffmann D. Toxic and carcinogenic agents in undiluted mainstream smoke and sidestream smoke of different types of cigarettes. *Carcinogenesis*, 1987; 8(5):729-31. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/3581431>.

Fletcher C and Peto R. The natural history of chronic airflow obstruction. *British Medical Journal*, 1977; 1(6077):1645-8. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1607732/>.

12.4.1 What is tobacco smoke?

Braun M, Klingelhofer D, Droge J, and Groneberg DA. Very high particulate matter levels by cigarettes from bangladesh and especially Nepal. *Nicotine & Tobacco Research*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35417555>

National Toxicology Program. Report on carcinogens, 15th edition. US Department of Health and Human Services, 2021. Available from:

<https://ntp.niehs.nih.gov/ntp/roc/content/profiles/tobaccorelatedexposures.pdf>.

Hu Y, Liu C, Xu Y, Yang J, and Pan Y. Identification of isobars and isomers in cigarette sidestream smoke in real time by synchrotron radiation photoionization mass spectrometry and multiple linear regression. *Analytical Chemistry*, 2021; 93(14):5718-26. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33797228>

Inaba Y, Uchiyama S, and Kunugita N. Spectrophotometric determination of ammonia levels in tobacco fillers of and sidestream smoke from different cigarette brands in Japan. *Environmental Health and Preventive Medicine*, 2018; 23(1):15. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29703135>

Diez-Izquierdo A, Cassanello-Penarroya P, Lidon-Moyano C, Matilla-Santander N, Balaguer A, et al. Update on thirdhand smoke: A comprehensive systematic review. *Environmental research*, 2018; 167:341-71. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30096604>

Wang H, Li X, Guo J, Peng B, Cui H, et al. Distribution of toxic chemicals in particles of various sizes from mainstream cigarette smoke. *Inhalation Toxicology*, 2016; 28(2):89-94. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26865272>

Noguchi M, Tanaka S, Watanabe K, and Yamasaki A. Correlation between odor concentration and volatile organic compounds (voc) composition of environmental Tobacco smoke (ETS). *Int J Environ Res Public Health*, 2016; 13(10). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27735848>

Li X, Lin B, Zhang H, Xie F, Ta N, et al. Cytotoxicity and mutagenicity of sidestream cigarette smoke particulate matter of different particle sizes. *Environmental Science and Pollution Research International*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26432262>

Daher N, Saleh R, Jaroudi E, Sheheitli H, Badr T, et al. Comparison of carcinogen, carbon monoxide, and ultrafine particle emissions from narghile waterpipe and cigarette smoking: Sidestream smoke measurements and assessment of second-hand smoke emission factors. *Atmospheric Environment*, 2010; 44(1):8-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20161525>

Polichetti G, Cocco S, Spinali A, Trimarco V, and Nunziata A. Effects of particulate matter (pm(10), pm(2.5) and pm(1)) on the cardiovascular system. *Toxicology*, 2009; 261(1–2):1–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19379789>

Moir D, Rickert WS, Levasseur G, Larose Y, Maertens R, et al. A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two machine smoking conditions. *Chemical Research in Toxicology*, 2008; 21(2):494-502. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18062674>

US Department of Health and Human Services. The health consequences of involuntary exposure to tobacco smoke: A report of the Surgeon General. Atlanta, Georgia: US Dept. of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2006. Available from: http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_2006/index.htm

Schick S and Glantz SA. Sidestream cigarette smoke toxicity increases with aging and exposure duration. *Tobacco Control*, 2006; 15(6):424–9. Available from: <http://tc.bmj.com/cgi/content/abstract/15/6/424>

Schick S and Glantz S. Philip Morris toxicological experiments with fresh sidestream smoke: More toxic than mainstream smoke. *Tobacco Control*, 2005; 14(6):396-404. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16319363>

Dobson R. Smoke from cigarette tip is more toxic than main inhaled smoke. *BMJ (Clinical Research Ed.)*, 2005; 331:1425. Available from: <http://www.bmj.com/cgi/content/extract/331/7530/1425-b>

Borgerding M and Klus H. Analysis of complex mixtures—cigarette smoke. *Experimental and Toxicologic Pathology*, 2005; 57 Suppl 1:43-73. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16092717>

International Agency for Research on Cancer Working Group on the Evaluation of Carcinogenic Risks to Humans. Tobacco smoke and involuntary smoking. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, Volume 83. Lyon: International Agency for Research on Cancer 2004. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Tobacco-Smoke-And-Involuntary-Smoking-2004>.

Kim YM, Harrad S, and Harrison RM. Concentrations and sources of VOCs in urban domestic and public microenvironments. *Environmental Science & Technology*, 2001; 35(6):997-1004. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11347947>

National Toxicology Program. Report on carcinogens, 9th edition. US Department of Health and Human Services, 1999. Available from: https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/tobacco_smoking1_2apps_508.pdf.

National Toxicology Program. Ntp report on carcinogens background document for tobacco smoking, final March 1999. Prepared for the October 30-31, 1997, Meeting of the Report on Carcinogens Subcommittee of the NTP Board of Scientific Counselors US Department of Health and Human Services, 1999. Available from: https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/tobacco_smoking1_2apps_508.pdf.

National Toxicology Program. Ntp report on carcinogens background document for environmental tobacco smoke. Prepared for the December 2-3, 1998, Meeting of the Report on Carcinogens Subcommittee of the NTP Board of Scientific Counselors US Department of Health and Human Services, 1998. Available from:

https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/environmentalts_noapps_508.pdf.

Heavner DL, Morgan WT, and Ogden MW. Determination of volatile organic compounds and ETS apportionment in 49 homes Environment International, 1995. Available from:

https://hero.epa.gov/hero/index.cfm/reference/details/reference_id/22045

Mohtashamipur E, Mohtashamipur A, Germann PG, Ernst H, Norpoth K, et al. Comparative carcinogenicity of cigarette mainstream and sidestream smoke condensates on the mouse skin. Journal of Cancer Research and Clinical Oncology, 1990; 116(6):604-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/2254379>

Brunnemann KD, Kagan MR, Cox JE, and Hoffmann D. Analysis of 1,3-butadiene and other selected gas-phase components in cigarette mainstream and sidestream smoke by gas chromatography-mass selective detection. Carcinogenesis, 1990; 11(10):1863-8. Available from:

<https://pubmed.ncbi.nlm.nih.gov/2208599/>

Vu-Duc T and Cong-Khanh H. Sidestream tobacco smoke constituents in indoor air modelled in an experimental chamber: Polycyclic aromatic hydrocarbons. Environment International, 1989; 15:57 - 64. Available from: <https://www.sciencedirect.com/science/article/pii/016041208990010X>

National Research Council (US) Committee on Passive Smoking, Environmental tobacco smoke: Measuring exposures and assessing health effects. Washington (DC): National Academies Press (US); 1986. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK219205/>

International Agency for Research on Cancer, Tobacco smoking. IARC monographs on the evaluation of carcinogen risk of chemicals to humans. Vol. 38 Vol. 38. Lyon: World Health Organization; 1986. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Tobacco-Smoking-1986>.

Woods JD and Harllee GC. Historical review of smoke pH data and sales trends for competitive brand filter cigarettes. RJ Reynolds, 10 May 1973 . Minnesota trial exhibit 12,337. 1973.

Bridge DP and Corn M. Contribution to the assessment of exposure of nonsmokers to air pollution from cigarette and cigar smoke in occupied spaces. Environmental research, 1972; 5(2):192-209. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/5036964>

12.4.1.1 Particles and gases in tobacco smoke

Mori, A, Ito, S, & Sekine, T. (2024). A revision of the multiple-path particle dosimetry model focusing on tobacco product aerosol dynamics. *Int J Numer Method Biomed Eng*, e3796. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38185887>

Zhang S, Wang Z, Zhang J, Guo D, and Chen Y. Inhalable cigarette-burning particles: Size-resolved chemical composition and mixing state. Environmental research, 2021:111790. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34329636>

Wang H, Li X, Guo J, Peng B, Cui H, et al. Distribution of toxic chemicals in particles of various sizes from mainstream cigarette smoke. *Inhalation Toxicology*, 2016; 28(2):89-94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26865272>

aher N, Saleh R, Jaroudi E, Sheheitli H, Badr T, et al. Comparison of carcinogen, carbon monoxide, and ultrafine particle emissions from narghile waterpipe and cigarette smoking: Sidestream smoke measurements and assessment of second-hand smoke emission factors. *Atmospheric Environment*, 2010; 44(1):8-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20161525>

Borgerding M and Klus H. Analysis of complex mixtures—cigarette smoke. *Experimental and Toxicologic Pathology*, 2005; 57 Suppl 1:43-73. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16092717>

International Agency for Research on Cancer, Tobacco smoking. IARC monographs on the evaluation of carcinogen risk of chemicals to humans. Vol. 38 Vol. 38. Lyon: World Health Organization; 1986. Available from: <https://publications.iarc.fr/Book-And-Report-Series/IARC-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Tobacco-Smoking-1986>.

Woods JD and Harllee GC. Historical review of smoke pH data and sales trends for competitive brand filter cigarettes. *RJ Reynolds*, 10 May 1973 . Minnesota trial exhibit 12,337. 1973.

12.4.1.2 Mainstream smoke, sidestream smoke and secondhand smoke

National Toxicology Program. Report on carcinogens, 15th edition. US Department of Health and Human Services, 2021. Available from: <https://ntp.niehs.nih.gov/ntp/roc/content/profiles/tobaccorelatedexposures.pdf>

Hu Y, Liu C, Xu Y, Yang J, and Pan Y. Identification of isobars and isomers in cigarette sidestream smoke in real time by synchrotron radiation photoionization mass spectrometry and multiple linear regression. *Analytical Chemistry*, 2021; 93(14):5718-26. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33797228>

Inaba Y, Uchiyama S, and Kunugita N. Spectrophotometric determination of ammonia levels in tobacco fillers of and sidestream smoke from different cigarette brands in Japan. *Environmental Health and Preventive Medicine*, 2018; 23(1):15. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29703135>

iez-Izquierdo A, Cassanello-Penarroya P, Lidon-Moyano C, Matilla-Santander N, Balaguer A, et al. Update on thirdhand smoke: A comprehensive systematic review. *Environmental research*, 2018; 167:341-71. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30096604>

oguchi M, Tanaka S, Watanabe K, and Yamasaki A. Correlation between odor concentration and volatile organic compounds (voc) composition of environmental Tobacco smoke (ETS). *Int J Environ Res Public Health*, 2016; 13(10). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27735848>

Li X, Lin B, Zhang H, Xie F, Ta N, et al. Cytotoxicity and mutagenicity of sidestream cigarette smoke particulate matter of different particle sizes. *Environmental Science and Pollution Research International*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26432262>

Daher N, Saleh R, Jaroudi E, Sheheitli H, Badr T, et al. Comparison of carcinogen, carbon monoxide, and ultrafine particle emissions from narghile waterpipe and cigarette smoking: Sidestream smoke measurements and assessment of second-hand smoke emission factors. *Atmospheric Environment*, 2010; 44(1):8-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20161525>

Polichetti G, Cocco S, Spinali A, Trimarco V, and Nunziata A. Effects of particulate matter (pm(10), pm(2.5) and pm(1)) on the cardiovascular system. *Toxicology*, 2009; 261(1–2):1–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19379789>

Moir D, Rickert WS, Levasseur G, Larose Y, Maertens R, et al. A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two machine smoking conditions. *Chemical Research in Toxicology*, 2008; 21(2):494-502. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18062674>

US Department of Health and Human Services. The health consequences of involuntary exposure to tobacco smoke: A report of the Surgeon General. Atlanta, Georgia: US Dept. of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2006. Available from: http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_2006/index.htm.

Schick S and Glantz SA. Sidestream cigarette smoke toxicity increases with aging and exposure duration. *Tobacco Control*, 2006; 15(6):424–9. Available from: <http://tc.bmj.com/cgi/content/abstract/15/6/424>

Schick S and Glantz S. Philip Morris toxicological experiments with fresh sidestream smoke: More toxic than mainstream smoke. *Tobacco Control*, 2005; 14(6):396-404. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16319363>

Dobson R. Smoke from cigarette tip is more toxic than main inhaled smoke. *BMJ (Clinical Research Ed.)*, 2005; 331:1425. Available from: <http://www.bmj.com/cgi/content/extract/331/7530/1425-b>

International Agency for Research on Cancer Working Group on the Evaluation of Carcinogenic Risks to Humans. Tobacco smoke and involuntary smoking. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, Volume 83. Lyon: International Agency for Research on Cancer 2004. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Tobacco-Smoke-And-Involuntary-Smoking-2004>.

Kim YM, Harrad S, and Harrison RM. Concentrations and sources of VOCs in urban domestic and public microenvironments. *Environmental Science & Technology*, 2001; 35(6):997-1004. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11347947>

National Toxicology Program. Report on carcinogens, 9th edition. US Department of Health and Human Services, 1999. Available from: https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/tobacco_smoking1_2apps_508.pdf.

National Toxicology Program. Ntp report on carcinogens background document for tobacco smoking, final March 1999. Prepared for the October 30-31, 1997, Meeting of the Report on Carcinogens Subcommittee of the NTP Board of Scientific Counselors US Department of Health and Human Services, 1999. Available from: https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/tobacco_smoking1_2apps_508.pdf.

National Toxicology Program. Ntp report on carcinogens background document for environmental tobacco smoke. Prepared for the December 2-3, 1998, Meeting of the Report on Carcinogens Subcommittee of the NTP Board of Scientific Counselors US Department of Health and Human Services, 1998. Available from: https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/environmentalts_noapps_508.pdf.

Heavner DL, Morgan WT, and Ogden MW. Determination of volatile organic compounds and ETS apportionment in 49 homes Environment International, 1995. Available from: https://hero.epa.gov/hero/index.cfm/reference/details/reference_id/22045

Mohtashamipur E, Mohtashamipur A, Germann PG, Ernst H, Norpoth K, et al. Comparative carcinogenicity of cigarette mainstream and sidestream smoke condensates on the mouse skin. *Journal of Cancer Research and Clinical Oncology*, 1990; 116(6):604-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/2254379>

Brunnemann KD, Kagan MR, Cox JE, and Hoffmann D. Analysis of 1,3-butadiene and other selected gas-phase components in cigarette mainstream and sidestream smoke by gas chromatography-mass selective detection. *Carcinogenesis*, 1990; 11(10):1863-8. Available from: <https://pubmed.ncbi.nlm.nih.gov/2208599/>

Vu-Duc T and Cong-Khanh H. Sidestream tobacco smoke constituents in indoor air modelled in an experimental chamber: Polycyclic aromatic hydrocarbons. *Environment International*, 1989; 15:57 - 64. Available from: <https://www.sciencedirect.com/science/article/pii/016041208990010X>

National Research Council (US) Committee on Passive Smoking, Environmental tobacco smoke: Measuring exposures and assessing health effects. Washington (DC): National Academies Press (US); 1986. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK219205/>

Bridge DP and Corn M. Contribution to the assessment of exposure of nonsmokers to air pollution from cigarette and cigar smoke in occupied spaces. *Environmental research*, 1972; 5(2):192-209. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/5036964>

12.4.2 What happens when tobacco products are burned?

Hall MG, Ribisl KM, and Brewer NT. Smokers' and nonsmokers' beliefs about harmful Tobacco constituents: Implications for FDA communication efforts. *Nicotine & Tobacco Research*, 2014; 16(3):343-50. Available from: <http://dx.doi.org/10.1093/ntr/ntt158>

Liu C, Feng S, van Heemst J, and McAdam K. New insights into the formation of volatile compounds in mainstream cigarette smoke. *Analytical and Bioanalytical Chemistry*, 2010; 396(5):1817-30. Available from: <http://www.springerlink.com/content/h0lj7116w543kk85/fulltext.html>

Kane D, Asgharian B, Price O, Rostami A, and Oldham M. Effect of smoking parameters on the particle size distribution and predicted airway deposition of mainstream cigarette smoke. *Inhalation Toxicology*, 2010; 22(3):199–209. Available from: <http://informahealthcare.com/doi/pdf/10.3109/08958370903161224>

Baker RR. Smoke generation inside a burning cigarette: Modifying combustion to develop cigarettes that may be less hazardous to health. *Progress in energy and Combustion Science*, 2006; 32:373-85. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S036012850600013X>

Ermala P and Holsti LR. On the burning temperatures of tobacco. *Cancer Research*, 1956; 16(6):490-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/13343119>

12.4.2.1 Pyrolysis

Wu, Z, Zhang, Q, Yu, H, Fu, L, Yang, Z, Lu, Y et al. (2024). Quantitative analysis of pyrolysis characteristics and chemical components of tobacco materials based on machine learning. *Front Chem*, 12, 1353745. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38380396>

Wei, H, Peng, Y, Huang, H, Fan, J, Xing, J, Luo, K et al. (2022). Toba-CPD: An Extended Chemical Percolation Devolatilization Model for Tobacco Pyrolysis. *ACS Omega*, 7(41), 36776-36785. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36278092>

Mu Y, Peng Y, Tang X, Ren J, Xing J, et al. Experimental and kinetic studies on Tobacco pyrolysis under a wide range of heating rates. *ACS Omega*, 2022; 7(1):1420-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35036803>

Chen YC, Chen KF, Lin JH, Huang SW, Chen HH, et al. The impact of pyrolysis temperature on physicochemical properties and pulmonary toxicity of tobacco stem micro-biochar. *Chemosphere*, 2021; 263:128349. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33297274>

Paschke M, Hutzler C, Henkler F, and Luch A. Oxidative and inert pyrolysis on-line coupled to gas chromatography with mass spectrometric detection: On the pyrolysis products of tobacco additives. *Int J Hyg Environ Health*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27622657>

Kibet JK, Khachatryan L, and Dellinger B. Phenols from pyrolysis and co-pyrolysis of tobacco biomass components. *Chemosphere*, 2015; 138:259-65. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26091866>

Boslaugh SE. Pyrolysis: Chemical reaction, in *Britannica*2014. Available from: <https://www.britannica.com/science/pyrolysis>.

Oja V, Hajaligol MR, and Waymack BE. The vaporization of semi-volatile compounds during tobacco pyrolysis. *Journal of Analytical and Applied Pyrolysis*, 2006; 76(1-2):117-23. Available from: <https://www.sciencedirect-com.ezproxy.une.edu.au/science/article/pii/S0165237005001646#bib17>

Baker RR. A review of pyrolysis studies to unravel reaction steps in burning tobacco. *Journal of Analytical and Applied Pyrolysis*, 1987; 11:555-73. Available from: <https://www.sciencedirect.com/science/article/abs/pii/0165237087850544>

Ermala P and Holsti LR. On the burning temperatures of tobacco. *Cancer Research*, 1956; 16(6):490-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/13343119>

12.4.2.2 Combustion

Moldoveanu SC. Interconversion of nicotine enantiomers during heating and implications for smoke from combustible cigarettes, heated tobacco products, and electronic cigarettes. *Chirality*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35088460>

Jebet A, Kibet J, Ombaka L, and Kinyanjui T. Surface bound radicals, char yield and particulate size from the burning of tobacco cigarette. *Chemistry Central Journal*, 2017; 11(1):79. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29086875>

McAdam K, Eldridge A, Fearon IM, Liu C, Manson A, et al. Influence of cigarette circumference on smoke chemistry, biological activity, and smoking behaviour. *Regulatory Toxicology and Pharmacology*, 2016; 82:111-26. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27634061>

Boslaugh SE. Pyrolysis: Chemical reaction, in *Britannica*2014. Available from: <https://www.britannica.com/science/pyrolysis>.

Kondratiev VN. Combustion: Chemical reaction, in *Britannica*2007. Available from: <https://www.britannica.com/science/combustion>.

Baker RR. Smoke generation inside a burning cigarette: Modifying combustion to develop cigarettes that may be less hazardous to health. *Progress in energy and Combustion Science*, 2006; 32:373-85. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S036012850600013X>

International Agency for Research on Cancer Working Group on the Evaluation of Carcinogenic Risks to Humans. Tobacco smoke and involuntary smoking. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, Volume 83. Lyon: International Agency for Research on Cancer 2004. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Tobacco-Smoke-And-Involuntary-Smoking-2004>.

12.4.2.3 Formation of smoke in cigarettes

Wei, J, Xiao, H, Wang, X, Zhao, H, Wang, X, Yao, S et al. (2023). Analysis of parametric instability of cigarettes based on computational fluid dynamics methods. *Heliyon*, 9(9), e19449. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37662803>

Gerlach G, Braun M, Droge J, and Groneberg DA. Do budget cigarettes emit more particles? An aerosol spectrometric comparison of particulate matter concentrations between private-label cigarettes and more expensive brand-name cigarettes. *Int J Environ Res Public Health*, 2022; 19(10). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35627457>

McAdam K, Eldridge A, Fearon IM, Liu C, Manson A, et al. Influence of cigarette circumference on smoke chemistry, biological activity, and smoking behaviour. *Regulatory Toxicology and Pharmacology*, 2016; 82:111-26. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27634061>

Shahab L, Hammond D, O'Connor R, Michael Cummings K, Borland R, et al. The reliability and validity of self-reported puffing behavior: Evidence from a cross-national study. *Nicotine & Tobacco Research*, 2008; 10(5):867–74. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200802027156>

12.4.2.4 Formation of smoke in other types of combusted tobacco products

Lee JW, Yang W, Kim YS, Kim Y, Yoo HS, et al. Exposure to secondhand smoke and a Tobacco-specific carcinogen in non-smokers. *Korean J Fam Med*, 2022; 43(2):117-24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35320897>

Vu AT, Hassink MD, Taylor KM, McGuigan M, Blasiolo A, et al. Volatile organic compounds in mainstream smoke of sixty domestic little cigar products. *Chemical Research in Toxicology*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33512154>

Jablonski JJ, Maines JH, Cheetham AG, and Gillman IG. Comparative levels of carbonyl delivery between mass-market cigars and cigarettes. *Regulatory Toxicology and Pharmacology*, 2019; 108:104453. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31473262>

Cartanya-Hueso A, Lidon-Moyano C, Fu M, Perez-Ortuno R, Ballbe M, et al. Comparison of tsnas concentration in saliva according to type of tobacco smoked. *Environmental research*, 2019; 172:73-80. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30771628>

Rosenberry ZR, Pickworth WB, and Koszowski B. Large cigars: Smoking topography and toxicant exposure. *Nicotine & Tobacco Research*, 2018; 20(2):183-91. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27798089>

Pickworth WB, Rosenberry ZR, Yi D, Pitts EN, Lord-Adem W, et al. Cigarillo and little cigar mainstream smoke constituents from replicated human smoking. *Chemical Research in Toxicology*, 2018; 31(4):251-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29582659>

Pickworth WB, Rosenberry ZR, and Koszowski B. Toxicant exposure from smoking a little cigar: Further support for product regulation. *Tobacco Control*, 2017; 26(3):269-76. Available from: <http://tobaccocontrol.bmj.com/content/tobaccocontrol/26/3/269.full.pdf>

Hamad SH, Johnson NM, Tefft ME, Brinkman MC, Gordon SM, et al. Little cigars vs 3R4F cigarette: Physical properties and HPHC yields. *Tobacco Regulatory Science*, 2017; 3(4):459-78. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29911130>

Ghosh A, Abdelwahab SH, Reeber SL, Reidel B, Marklew AJ, et al. Little cigars are more toxic than cigarettes and uniquely change the airway gene and protein expression. *Sci Rep*, 2017; 7:46239. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28447619>

Dethloff O, Mueller C, Cahours X, and Colard S. Cigar burning under different smoking intensities and effects on emissions. *Regulatory Toxicology and Pharmacology*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29074275>

Klupinski TP, Strozier ED, Friedenber DA, Brinkman MC, Gordon S, et al. Identification of new and distinctive exposures from little cigars. *Chemical Research in Toxicology*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26605856>

Piade JJ, Roemer E, Dempsey R, Hornig G, Deger Evans A, et al. Toxicological assessment of kretek cigarettes: Part 2: Kretek and American-blended cigarettes, smoke chemistry and in vitro toxicity. *Regulatory Toxicology and Pharmacology*, 2014; 70 Suppl 1:S15-25. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25497993>

Koszowski B, Rosenberry ZR, Viray LC, Potts JL, and Pickworth WB. Make your own cigarettes: Toxicant exposure, smoking topography, and subjective effects. *Cancer Epidemiology Biomarkers & Prevention*, 2014. Available from: <http://cebp.aacrjournals.org/content/cebp/early/2014/06/12/1055-9965.EPI-14-0280.full.pdf>

Ahamad T and Alshehri SM. Tg-ftir-ms (evolved gas analysis) of bidi tobacco powder during combustion and pyrolysis. *Journal of Hazardous Materials*, 2012; 199-200:200-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22119196>

Shahab L, West R, and McNeill A. A comparison of exposure to carcinogens among roll-your-own and factory-made cigarette smokers. *Addiction Biology*, 2009; 14(3):315-20. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19523045>

Gupta PC and Asma S. Bidi smoking and public health. Nee Delhi: Ministry of Health and Family Welfare, Government of India, 2008. Available from: https://www.healis.org/pdf/special-report/Bidi_smoking_and_public_health.pdf.

International Agency for Research on Cancer Working Group on the Evaluation of Carcinogenic Risks to Humans. Tobacco smoke and involuntary smoking. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, Volume 83. Lyon: International Agency for Research on Cancer 2004. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Tobacco-Smoke-And-Involuntary-Smoking-2004>.

Rahman M and Fukui T. Bidi smoking and health. *Public Health*, 2000; 114(2):123-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/10800151>

Klepeis NE, Ott WR, and Repace JL. The effect of cigar smoking on indoor levels of carbon monoxide and particles. *Journal of Exposure Analysis and Environmental Epidemiology*, 1999; 9(6):622-35. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/10638847>

Hoffman D and Hoffman I. Cigar smoke: Chemistry and toxicology. *Smoking and tobacco control monograph No. 9*, 1998. Available from: https://cancercontrol.cancer.gov/sites/default/files/2020-06/m9_3.pdf.

Ohshima H, Nair J, Bourgade MC, Friesen M, Garren L, et al. Identification and occurrence of two new n-nitrosamino acids in tobacco products: 3-(N-nitroso-N-methylamino)propionic acid and 4-(N-nitroso-N-methylamino)butyric acid. *Cancer Letters*, 1985; 26(2):153-62. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/3978605>

Ermala P and Holsti LR. On the burning temperatures of tobacco. *Cancer Research*, 1956; 16(6):490-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/13343119>

12.4.3 Chemicals that are emitted from combusted tobacco products

El-Hellani, A, Adeniji, A, Erythropel, HC, Wang, Q, Lamb, T, Mikheev, VB et al. (2024). Comparison of emissions across tobacco products: A slippery slope in tobacco control. *Tob Induc Dis*, 22. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38560551>

Wendt, CH, Bowler, RP, Demorest, C, Hastie, A, Labaki, WW, Chen, M et al. (2023). Levels of Urinary Mercapturic Acids of Acrolein, Methacrolein, Crotonaldehyde, and Methyl Vinyl Ketone in

Relationship to Chronic Obstructive Pulmonary Disease in Cigarette Smokers of the Subpopulations and Intermediate Outcome Measures in COPD Study (SPIROMICS). *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37725788>

Zeng, T, Liu, Y, Jiang, Y, Zhang, L, Zhang, Y, Zhao, L et al. (2023). Advanced Materials Design for Adsorption of Toxic Substances in Cigarette Smoke. *Adv Sci (Weinh)*, e2301834. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37211707>

Maiyo AK, Kibet JK, and Kengara FO. A review of the characteristic properties of selected tobacco chemicals and their associated etiological risks. *Reviews on Environmental Health*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35538694>

Lim DH, Son YS, Kim YH, Kukkar D, and Kim KH. Volatile organic compounds released in the mainstream smoke of flavor capsule cigarettes. *Environmental research*, 2022;112866. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35134376>

Li Y and Hecht SS. Carcinogenic components of tobacco and tobacco smoke: A 2022 update. *Food and Chemical Toxicology*, 2022; 165:113179. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35643228>

Jung CC, Syu ZH, Su HJ, Lian PY, and Chen NT. Stable c and n isotopes of pm2.5 and size-segregated particles emitted from incense stick and cigarette burning. *Environmental research*, 2022; 212(Pt B):113346. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35461851>

Deng B, Wang Y, Huang H, Duan X, and Liu A. Effects of inhalation frequency on inhalation/exposure dose of hazardous nanoparticles and toxic gases during cigarette smoking. *Ecotoxicol Environ Saf*, 2022; 240:113709. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35653970>

Amoros-Perez A, Cano-Casanova L, Roman-Martinez MDC, and Lillo-Rodenas MA. Solid matter and soluble compounds collected from cigarette smoke and heated tobacco product aerosol using a laboratory designed puffing setup. *Environmental research*, 2022; 206:112619. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34971599>

National Toxicology Program. Report on carcinogens, 15th edition. US Department of Health and Human Services, 2021. Available from: <https://ntp.niehs.nih.gov/ntp/roc/content/profiles/tobaccorelatedexposures.pdf>.

Mittelstaedt RA, Shaddock JG, Bhalli JA, Guo X, Li Y, et al. Differentiating between micronucleus dose-responses induced by whole cigarette smoke solutions with benchmark dose potency ranking. *Mutation Research*, 2021; 866:503351. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33985695>

Wei M, Zheng F, Song X, Li R, Pan X, et al. Comprehensive analysis of volatile compounds in mouthpiece cigarette adhesive by coupling headspace with gas chromatography-mass spectrometry. *Journal of AOAC International*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33252696>

Pack EC, Kim HS, Lee SH, Koo YJ, Jang DY, et al. Survey of characteristics of exposure to mainstream cigarette smoke using discarded cigarette butts from Korean smokers. *Environmental research*, 2020; 185:109434. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32276166>

Graves BM, Johnson TJ, Nishida RT, Dias RP, Savareear B, et al. Comprehensive characterization of mainstream marijuana and tobacco smoke. *Sci Rep*, 2020; 10(1):7160. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32345986>

World Health Organization. Report on the scientific basis of tobacco product regulation: Seventh report of a WHO study group. Geneva: WHO, 2019. Available from: <https://www.who.int/publications/i/item/who-study-group-on-tobacco-product-regulation-report-on-the-scientific-basis-of-tobacco-product-regulation-seventh-report-of-a-who-study-group>.

St Helen G, Benowitz NL, Ko J, Jacob P, Gregorich SE, et al. Differences in exposure to toxic and/or carcinogenic volatile organic compounds between black and white cigarette smokers. *Journal of Exposure Science & Environmental Epidemiology*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31406274>

Schwanz TG, Bokowski LVV, Marcelo MCA, Jandrey AC, Dias JC, et al. Analysis of chemosensory markers in cigarette smoke from different tobacco varieties by gcxgc-tofms and chemometrics. *Talanta*, 2019; 202:74-89. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31171230>

Pennings JLA, Cremers J, Becker MJA, Klerx WNM, and Talhout R. Aldehyde and voc yields in commercial cigarette mainstream smoke are mutually related and depend on the sugar and humectant content in tobacco. *Nicotine & Tobacco Research*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31679033>

Jeong M, Noar SM, Zhang D, Mendel JR, Agans RP, et al. Public understanding of cigarette smoke chemicals: Longitudinal study of US adults and adolescents. *Nicotine & Tobacco Research*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30852611>

Hecht SS and DeMarini DM. Tobacco smoke and its constituents, in Tumour site concordance and mechanisms of carcinogenesis. Baan RA, Stewart BW, and Straif K, Editors. Lyon (FR): 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33979081>.

Cai B, Li Z, Wang R, Geng Z, Shi Y, et al. Emission level of seven mainstream smoke toxicants from cigarette with variable tobacco leaf constituents. *Regulatory Toxicology and Pharmacology*, 2019; 103:181-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30710578>

Zumbado M, Luzardo OP, Rodriguez-Hernandez A, Boada LD, and Henriquez-Hernandez LA. Differential exposure to 33 toxic elements through cigarette smoking, based on the type of tobacco and rolling paper used. *Environmental research*, 2018; 169:368-76. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30513508>

World Health Organization. Tobacco product regulation: Basic handbook. Geneva: WHO, 2018. Available from: <https://www.who.int/publications/i/item/tobacco-product-regulation-basic-handbook>.

Tobacco laboratory Network. Determination of volatile organics in mainstream cigarette smoke under ISO and intense smoking conditions. SOP 09. Geneva: WHO, 2018. Available from: <https://apps.who.int/iris/bitstream/handle/10665/275344/9789241514774-eng.pdf>.

Pack EC, Jang DY, Kim HS, Lee SH, Kim HY, et al. Mixture risk assessment of selected mainstream cigarette smoke constituents generated from low-yield cigarettes in south Korean smokers.

- Regulatory Toxicology and Pharmacology, 2018; 94:152-62. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29408505>
- Mendel JR, Baig SA, Hall MG, Jeong M, Byron MJ, et al. Brand switching and toxic chemicals in cigarette smoke: A national study. PloS One, 2018; 13(1):e0189928. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29324749>
- Marcilla A, Beltrán MI, Gómez-Siurana A, Martínez Castellanos I, and Berenguer Muñoz D. Comparative study of the composition of the mainstream smoke of five brands factory-made-cigarettes and their counterpart roll-your-own tobaccos. Universal Journal of Management, 2018; 6(1):12-23. Available from: <http://hdl.handle.net/10045/73588>
- Lee PN. Tar level of cigarettes smoked and risk of smoking-related diseases. Inhalation Toxicology, 2018; 30(1):5-18. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29488428>
- Kim YH, An YJ, Jo S, Lee SH, Lee SJ, et al. Comparison of volatile organic compounds between cigarette smoke condensate (csc) and extract (cse) samples. Environmental Health and Toxicology, 2018; 33(3):e2018012-0. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30286588>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6182245/pdf/eht-33-3-e2018012.pdf>
- Goel R, Bitzer ZT, Reilly SM, Foulds J, Muscat J, et al. Influence of smoking puff parameters and Tobacco varieties on free radicals yields in cigarette mainstream smoke. Chemical Research in Toxicology, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29701955>
- Goel R, Bitzer Z, Reilly SM, Trushin N, Foulds J, et al. Variation in free radical yields from U.S. Marketed cigarettes. Chemical Research in Toxicology, 2017; 30(4):1038-45. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28269983>
- Ding YS, Richter P, Hearn B, Zhang L, Bravo R, et al. Chemical characterization of mainstream smoke from spectrum variable nicotine research cigarettes. Tob Regul Sci, 2017; 3(1):81-94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28220149>
- Brewer NT, Morgan JC, Baig SA, Mendel JR, Boynton MH, et al. Public understanding of cigarette smoke constituents: Three US surveys. Tobacco Control, 2017; 26(5):592-9. Available from: <http://tobaccocontrol.bmj.com/content/tobaccocontrol/26/5/592.full.pdf>
- Wang H, Li X, Guo J, Peng B, Cui H, et al. Distribution of toxic chemicals in particles of various sizes from mainstream cigarette smoke. Inhalation Toxicology, 2016; 28(2):89-94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26865272>
- Pazo DY, Moliere F, Sampson MM, Reese CM, Agnew-Heard KA, et al. Mainstream smoke levels of volatile organic compounds in 50 US domestic cigarette brands smoked with the ISO and canadian intense protocols. Nicotine & Tobacco Research, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27113015>
- Li X. In vitro toxicity testing of cigarette smoke based on the air-liquid interface exposure: A review. Toxicology In Vitro, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27470133>
- Kant N, Muller R, Braun M, Gerber A, and Groneberg D. Particulate matter in second-hand smoke emitted from different cigarette sizes and types of the brand vogue mainly smoked by women. Int J

Environ Res Public Health, 2016; 13(8). Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27509517>

Horiyama S, Kunitomo M, Yoshikawa N, and Nakamura K. Mass spectrometric approaches to the identification of potential ingredients in cigarette smoke causing cytotoxicity. *Biol Pharm Bull*, 2016; 39(6):903-8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27251491>

Alexandrov LB, Ju YS, Haase K, Van Loo P, Martincorena I, et al. Mutational signatures associated with tobacco smoking in human cancer. *Science*, 2016; 354(6312):618-22. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27811275>

Uchiyama S, Hayashida H, Izu R, Inaba Y, Nakagome H, et al. Determination of nicotine, tar, volatile organic compounds and carbonyls in mainstream cigarette smoke using a glass filter and a sorbent cartridge followed by the two-phase/one-pot elution method with carbon disulfide and methanol. *Journal of Chromatography A*, 2015; 1426:48-55. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26653840>

Morrissey H, Ball P, Boland M, Hefler M, and Thomas DP. Constituents of smoke from cigarettes made from diverted nicotine replacement therapy patches. *Drug and Alcohol Review*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26094737>

Eldridge A, Betson TR, Gama MV, and McAdam K. Variation in tobacco and mainstream smoke toxicant yields from selected commercial cigarette products. *Regulatory Toxicology and Pharmacology*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25620723>

Camacho OM, Eldridge A, Proctor CJ, and McAdam K. Empirical characterisation of ranges of mainstream smoke toxicant yields from contemporary cigarette products using quantile regression methodology. *Regulatory Toxicology and Pharmacology*, 2015; 72(3):458-72. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26021184>

Benowitz NL, Gan Q, Goniewicz ML, Lu W, Xu J, et al. Different profiles of carcinogen exposure in chinese compared with US cigarette smokers. *Tobacco Control*, 2015; 24(e4):e258-e63. Available from: <http://tobaccocontrol.bmj.com/content/24/e4/e258.abstract>

Schneller LM, Zwierzchowski BA, Caruso RV, Li Q, Yuan J, et al. Changes in tar yields and cigarette design in samples of chinese cigarettes, 2009 and 2012. *Tobacco Control*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25352560>

Oldham MJ, DeSoi DJ, Rimmer LT, Wagner KA, and Morton MJ. Insights from analysis for harmful and potentially harmful constituents (hphcs) in tobacco products. *Regulatory Toxicology and Pharmacology*, 2014; 70(1):138-48. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24973503>

Marcilla A, Beltran MI, Gomez-Siurana A, Berenguer D, and Martinez-Castellanos I. Comparison between the mainstream smoke of eleven RYO tobacco brands and the reference tobacco 3R4F. *Toxicology Reports*, 2014; 1:122-36.

Comer DM, Elborn JS, and Ennis M. Inflammatory and cytotoxic effects of acrolein, nicotine, acetaldehyde and cigarette smoke extract on human nasal epithelial cells. *BMC Pulm Med*, 2014; 14:32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24581246>

Talhout R, Schulz, T., Florek, E., van Benthem, J., Wester, P., Opperhuizen, A. Hazardous compounds in tobacco smoke. *Int J Environ Res Public Health*, 2011; 8:613-28.

US Department of Health and Human Services. A report of the Surgeon General: How tobacco smoke causes disease. US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2010. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK53017/>.

Nelson P, Chen P, Dixon M, and Steichen T. A survey of mouth level exposure to cigarette smoke in the United States. *Regulatory Toxicology and Pharmacology*, 2010; [Epub ahead of print]. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20937343

Roemer E, Ottmueller T, Zenzen V, Wittke S, Radtke F, et al. Cytotoxicity, mutagenicity, and tumorigenicity of mainstream smoke from three reference cigarettes machine-smoked to the same yields of total particulate matter per cigarette. *Food and Chemical Toxicology*, 2009; 47(8):1810–8. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19447158

Pickett G, Seagrave J, Boggs S, Polzin G, Richter P, et al. Effects of 10 cigarette smoke condensates on primary human airway epithelial cells by comparative gene and cytokine expression studies. *Toxicological Sciences*, 2009; 114(1):79–89. Available from: <http://toxsci.oxfordjournals.org/content/114/1/79.long>

Ejaz S, Insan-ud-din, Ashraf M, Nawaz M, Lim C, et al. Cigarette smoke condensate and total particulate matter severely disrupts physiological angiogenesis. *Food and Chemical Toxicology*, 2009; 47(3):601–14. Available from: http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6T6P-4V75YXF-6&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_acct=C000050221&_version=1&_urlVersion=0&_userid=10&md5=0c71936283973d98f85866d7db1edbc3

Hammond D and O'Connor RJ. Constituents in tobacco and smoke emissions from canadian cigarettes. *Tobacco Control*, 2008; 17(suppl. 1):i24-31. Available from: http://tobaccocontrol.bmj.com/cgi/content/abstract/17/Suppl_1/i24

Scientific Advisory Committee on Tobacco. The scientific basis of tobacco product regulation: Report of a WHO study group. Technical Report Series, no. 945. Geneva: World Health Organization, 2007. Available from: <https://escholarship.org/uc/item/4bx2m81n>

King B, Borland R, and Fowles J. Mainstream smoke emissions of Australian and canadian cigarettes. *Nicotine & Tobacco Research*, 2007; 9(8):835–44. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200701485109>

van Amsterdam J, Talhout R, Vleeming W, and Opperhuizen A. Contribution of monoamine oxidase (mao) inhibition to tobacco and alcohol addiction. *Life Sciences*, 2006; 79:1969-73.

Laugesen M and Fowles J. Marlboro ultrasmooth: A potentially reduced exposure cigarette? *Tobacco Control*, 2006; 15(6):430–5. Available from: <http://tc.bmj.com/cgi/content/abstract/15/6/430>

Hammond D, Collishaw NE, and Callard C. Secret science: Tobacco industry research on smoking behaviour and cigarette toxicity. *Lancet*, 2006; 367(9512):781-7. Available from: <https://pubmed.ncbi.nlm.nih.gov/16517278/>

Hecht SS, Murphy SE, Carmella SG, Li S, Jensen J, et al. Similar uptake of lung carcinogens by smokers of regular, light, and ultralight cigarettes. *Cancer Epidemiology, Biomarkers & Prevention*, 2005; 14(3):693-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15767351>

US Department of Health and Human Services, The health consequences of smoking: A report of the Surgeon General. Atlanta, Georgia: US Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health 2004. Available from: http://www.cdc.gov/tobacco/data_statistics/sgr/index.htm.

Djordjevic M, Stellman S, and Zang E. Doses of nicotine and lung carcinogens delivered to cigarette smokers. *Journal of National Cancer Institute*, 2000; 92(2):106–11. Available from: <http://jnci.oxfordjournals.org/cgi/content/full/92/2/106>

National Toxicology Program. Report on carcinogens, 9th edition. US Department of Health and Human Services, 1999. Available from: https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/tobacco_smoking1_2apps_508.pdf.

National Toxicology Program. Ntp report on carcinogens background document for tobacco smoking, final March 1999. Prepared for the October 30-31, 1997, Meeting of the Report on Carcinogens Subcommittee of the NTP Board of Scientific Counselors US Department of Health and Human Services, 1999. Available from: https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/tobacco_smoking1_2apps_508.pdf.

Meier WM and Siegman K. Significant reduction of carcinogenic compounds in tobacco smoke by the use of zeolite catalysts. *Microporous and Mesoporous Materials*, 1999; 33(1–3):307–10. Available from: http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6TH4-3YYT7PS-W&_user=559483&_rdoc=1&_fmt=&_orig=search&_sort=d&view=c&_version=1&_urlVersion=0&_urlserid=559483&md5=7dbcecb077c431f161a0b18fe06bbc40

National Toxicology Program. Ntp report on carcinogens background document for environmental tobacco smoke. Prepared for the December 2-3, 1998, Meeting of the Report on Carcinogens Subcommittee of the NTP Board of Scientific Counselors US Department of Health and Human Services, 1998. Available from: https://ntp.niehs.nih.gov/ntp/newhomeroc/other_background/environmentalts_noapps_508.pdf.

Smith CJ, Livingston SD, and Doolittle DJ. An international literature survey of “IARC group I carcinogens” reported in mainstream cigarette smoke. *Food and Chemical Toxicology*, 1997; 35(10-11):1107-30. Available from: <https://pubmed.ncbi.nlm.nih.gov/9463546/>

Hoffmann D, Djordjevic, MV and Brunnemann, KD. Changes in cigarette design and composition over time and how they influence the yields of smoke constituents., in *The FTC cigarette test method for determining tar, nicotine, and carbon monoxide yields of U.S. Cigarettes*. Smoking and Tobacco control monograph 7. Bethesda, MD: U.S. Department of Health and Human Services, Public Health

Service, National Institutes of Health.; 1996. p 15-37 Available from:
<https://cancercontrol.cancer.gov/brp/tcrb/monographs/monograph-07>.

Harris JE. Cigarette smoke components and disease: Cigarette smoke is more than a triad of tar, nicotine, and carbon monoxide, in The FTC cigarette test method for determining tar, nicotine and carbon monoxide yields of US cigarettes. Report of the nci expert committee. Usdhhs. 1996. Available from: cancercontrol.cancer.gov/tcrb/monographs/7/m7_5.pdf.

Djordjevic L and Hoffmann D. A comparison of selected components in the mainstream smoke of the leading US and Japanese cigarettes. in CORESTA. 1996.

International Agency for Research on Cancer. IARC monographs on the evaluation of carcinogenic risks to humans: Some industrial chemicals. 60 Lyon, France: IARC, 1994. Available from:
<https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Some-Industrial-Chemicals-1994>.

Hoffmann D. Analysis of toxic smoke constituents. Toxicity testing plan. Table 3: US Consumer Product Safety Commission and US Department of Health and Human Services, 1993.

Prignot J. Quantification and chemical markers of tobacco-exposure. European Respiratory Society, 1987; 70(1):1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/3545884>

Bradford JA, Harlan WR, and Hanmer HR. Nature of cigaret smoke. Industrial and engineering chemistry, 1936; 28(7):836 - 9. Available from: <https://pubs-acsc-org.ezproxy.une.edu.au/doi/abs/10.1021/ie50319a020>

US Department of Health and Human Services. Cigarette smoke components and disease: Cigarette smoke is more than a triad of tar, nicotine, and carbon monoxide. The FTC cigarette test method for determining tar, nicotine and carbon monoxide yields of US cigarettes. Report of the nci expert committee. US Department of Health and Human Services, Available from:
<http://tobaccodocuments.org/rjr/515648348-8640.html>.

12.4.3.1 Nicotine and other alkaloids

Cao, Y, Liu, X, Hu, Z, Li, J, Chen, X, Xiong, Y et al. (2024). Assessing nicotine pharmacokinetics of new generation tobacco products and conventional cigarettes: a systematic review and meta-analysis. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39162577>

Xue, F, Yang, J, Luo, C, Li, D, Shi, G, Song, G, & Li, Y. (2023). Metagenomic insight into the biodegradation of biomass and alkaloids in the aging process of cigar. *Bioresour Bioprocess*, 10(1), 45. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38647787>

Chu, M, Deng, J, Hu, H, Wang, R, Li, D, Chen, Z et al. (2023). Nicotine Transport across Calu-3 Cell Monolayer: Effect of Nicotine Salts and Flavored E-liquids. *Drug Dev Ind Pharm*, 1-27. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37751149>

Salam, S, El-Hajj Moussa, F, El-Hage, R, El-Hellani, A, & Aoun Saliba, N. (2023). A Systematic Review of Analytical Methods for the Separation of Nicotine Enantiomers and Evaluation of Nicotine Sources. *Chem Res Toxicol*, 36(3), 334-341. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36897818>

Rehan, M, Zargar, UR, Sheikh, IA, Alharthy, SA, Almashjary, MN, Abuzenadah, AM, & Beg, MA. (2022). Potential Disruption of Systemic Hormone Transport by Tobacco Alkaloids Using Computational Approaches. *Toxics*, 10(12) Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36548560>

Liu, G, Wang, R, Chen, H, Wu, P, Fu, Y, Li, K et al. (2022). Non-nicotine constituents in cigarette smoke extract enhance nicotine addiction through monoamine oxidase A inhibition. *Front Neurosci*, 16, 1058254. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36507317>

Guo, Q, Qin, Y, Pan, L, Xie, F, Liu, S, Sun, X et al. (2022). Accuracy improvement of determination of seven minor tobacco alkaloids in mainstream cigarette smoke using analyte protectants by gas chromatography-mass spectrometry. *J Chromatogr A*, 1684, 463537. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36240707>

Thasweer, AM, Renuka Devi, P, & Thirunavukkarasu, V. (2022). Molecular docking and dynamic simulation studies of alpha4beta2 and alpha7 nicotinic acetylcholine receptors with tobacco smoke constituents nicotine, NNK and NNN. *J Biomol Struct Dyn*, 1-10. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36270967>

Wilhelm J, Mishina E, Viray L, Paredes A, and Pickworth WB. The pH of smokeless tobacco determines nicotine buccal absorption: Results of a randomized crossover trial. *Clinical Pharmacology and Therapeutics*, 2022; 111(5):1066-74. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34826137>

Moldoveanu SC. Interconversion of nicotine enantiomers during heating and implications for smoke from combustible cigarettes, heated tobacco products, and electronic cigarettes. *Chirality*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35088460>

Li L, Zhang H, Wen J, Shen Y, Li D, et al. Direct determination of free nicotine content in Tobacco. *ACS Omega*, 2022; 7(27):23061-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35847304>

Qamar W, Altamimi MA, Rehman MU, Ali N, Imam F, et al. Toxicological interaction between tobacco smoke toxicants cadmium and nicotine: An in-vitro investigation. *Saudi Journal of Biological Sciences*, 2021; 28(8):4201-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34354400>

hang YJ, Muthukumaran RB, Chen JL, Chang HY, Hung YC, et al. Simultaneous determination of areca nut- and tobacco-specific alkaloids in saliva by lc-ms/ms: Distribution and transformation of alkaloids in oral cavity. *J Hazard Mater*, 2021; 426:128116. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34968842>

Tanner NT, Thomas NA, Ward R, Rojewski A, Gebregziabher M, et al. Association of cigarette type and nicotine dependence in patients presenting for lung cancer screening. *Chest*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32603713>

Miller JH, Danielson T, Pithawalla YB, Brown AP, Wilkinson C, et al. Method development and validation of dissolution testing for nicotine release from smokeless tobacco products using flow-

through cell apparatus and uplc-pda. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences*, 2020; 1141:122012. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32065955>

Li C, Li E, Ma M, Liu X, You J, et al. Simultaneous determination of six alkaloids in tobacco and tobacco products by direct analysis of real-time triple quadrupole mass spectrometry with a modified pretreatment method. *Journal of Separation Science*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32034866>

Lei X, Goel R, Sun D, Bhangu G, Bitzer ZT, et al. Free radical and nicotine yields in mainstream smoke of chinese marketed cigarettes: Variation with smoking regimens and cigarette brands. *Chemical Research in Toxicology*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32363856>

uiper N, Coats EM, Crawford TN, Gammon DG, Loomis B, et al. Trends in manufacturer-reported nicotine yields in cigarettes sold in the United States, 2013–2016. *Prev Chronic Dis*, 2020; 17:E148. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33241990>

Jing Y, Ning S, Guan Y, Cao M, Li J, et al. Electrochemical determination of nicotine in Tobacco products based on biosynthesized gold nanoparticles. *Front Chem*, 2020; 8:593070. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33195097>

Ebajemito JK, McEwan M, Gale N, Camacho OM, Hardie G, et al. A randomised controlled single-centre open-label pharmacokinetic study to examine various approaches of nicotine delivery using electronic cigarettes. *Sci Rep*, 2020; 10(1):19980. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33235307>

Salman R, Talih S, El-Hage R, Haddad C, Karaoghlanian N, et al. Free-base and total nicotine, reactive oxygen species, and carbonyl emissions from iqos, a heated tobacco product. *Nicotine & Tobacco Research*, 2019; 21(9):1285-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30476301>

Lin B, Chen J, Zeng Y, Li L, Qiu B, et al. A facile approach for on-site evaluation of nicotine in tobacco and environmental tobacco smoke. *ACS Sens*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31250643>

Zhang X, Wang R, Zhang L, Ruan Y, Wang W, et al. Simultaneous determination of tobacco minor alkaloids and tobacco-specific nitrosamines in mainstream smoke by dispersive solid-phase extraction coupled with ultra-performance liquid chromatography/tandem orbitrap mass spectrometry. *Rapid Communications in Mass Spectrometry*, 2018; 32(20):1791-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29964303>

Ng HW, Leggett C, Sakkiah S, Pan B, Ye H, et al. Competitive docking model for prediction of the human nicotinic acetylcholine receptor alpha7 binding of tobacco constituents. *Oncotarget*, 2018; 9(24):16899-916. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29682193>

Hellinghausen G, Roy D, Wang Y, Lee JT, Lopez DA, et al. A comprehensive methodology for the chiral separation of 40 tobacco alkaloids and their carcinogenic e/z-(r,s)-tobacco-specific nitrosamine metabolites. *Talanta*, 2018; 181:132-41. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29426492>

Zuo Y, Garg PK, Nazih R, Garg S, Rose JE, et al. A programmable smoke delivery device for pet imaging with cigarettes containing 11c-nicotine. *J Neurosci Methods*, 2017; 283:55-61. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28347784>

Tobacco laboratory Network. Determination of nicotine and carbon monoxide in mainstream cigarette smoke under intense smoking conditions. SOP 10. Geneva: WHO, 2016. Available from: <https://www.who.int/publications/i/item/standard-operating-procedure-for-determination-of-nicotine-and-carbon-monoxide-in-mainstream-cigarette-smoke-under-intense-smoking-conditions>.

Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). Final opinion on additives used in tobacco products. European Commission, Health & Food Safety, Directorate C: Public Health 2016. Available from:

http://ec.europa.eu/health/scientific_committees/emerging/docs/scenihr_o_051.pdf.

Perkins KA, Kunkle N, Michael VC, Karelitz JL, and Donny EC. Assessing discrimination of nicotine in humans via cigarette smoking. *Nicotine & Tobacco Research*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27013335>

Kurgat C, Kibet J, and Cheplogoi P. Molecular modeling of major tobacco alkaloids in mainstream cigarette smoke. *Chemistry Central Journal*, 2016; 10:43. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27429644>

Kibet J, Kurgat C, Limo S, Rono N, and Bosire J. Kinetic modeling of nicotine in mainstream cigarette smoking. *Chemistry Central Journal*, 2016; 10:60. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27790285>

Redner R, White TJ, Bunn JY, and Higgins ST. Use of high nicotine/tar yield (full-flavor) cigarettes and risk for nicotine dependence in nationally representative samples of U.S. Smokers. *Nicotine & Tobacco Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26547061>

Lin YH, Feng CH, Wang SW, Ko PY, Lee MH, et al. Determination of nicotine in Tobacco by chemometric optimization and cation-selective exhaustive injection in combination with sweeping-micellar electrokinetic chromatography. *Journal of Analytical Methods in Chemistry*, 2015; 2015:869719. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26101695>

Jarvis MJ, Giovino GA, O'Connor RJ, Kozlowski LT, and Bernert JT. Variation in nicotine intake in U.S. Cigarette smokers over the past 25 years: Evidence from nhanes surveys. *Nicotine & Tobacco Research*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25063772>

van Amsterdam J, Sleijffers A, van Spiegel P, Blom R, Witte M, et al. Effect of ammonia in cigarette tobacco on nicotine absorption in human smokers. *Food and Chemical Toxicology*, 2011; 49(12):3025-30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22001171>

St Charles F, Cook C, and Clayton P. The linear relationship between cigarette tar and nicotine yields: Regulatory implications for smoke constituent ratios. *Regulatory Toxicology and Pharmacology*, 2011; [Epub ahead of print]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21216263>

Lauterbach J, Bao M, Joza P, and Rickert W. Free-base nicotine in tobacco products. Part i. Determination of free-base nicotine in the particulate phase of mainstream cigarette smoke and the relevance of these findings to product design parameters. *Regulatory Toxicology and Pharmacology*, 2010; 58(1):45-63. Available from: <http://www.sciencedirect.com/science/journal/02732300>

Nishioka T, Guo J, Yamamoto D, Chen L, Huppi P, et al. Nicotine, through upregulating pro-survival signaling, cooperates with NNK to promote transformation. *Journal of Cellular Biochemistry*, 2009; 109(1):152–61. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/jcb.22392/full>

Gowadia N, Oldham M, and Dunn-Rankin D. Particle size distribution of nicotine in mainstream smoke from 2r4f, Marlboro medium, and quest1 cigarettes under different puffing regimens. *Inhalation Toxicology*, 2009; 21(5):435–46. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19496699

Benowitz NL, Hukkanen J, and Jacob P, 3rd. Nicotine chemistry, metabolism, kinetics and biomarkers. *Handbook of Experimental Pharmacology*, 2009; 192(192):29-60. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19184645>

Stevenson T and Proctor RN. The secret and soul of Marlboro: Phillip Morris and the origins, spread, and denial of nicotine freebasing. *American Journal of Public Health*, 2008; 98(7):1184-94. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18511721>

Ferris Wayne G, Connolly GN, and Henningfield JE. Brand differences of free-base nicotine delivery in cigarette smoke: The view of the tobacco industry documents. *Tobacco Control*, 2006; 15(3):189-98. Available from: <http://tc.bmjournals.com/cgi/content/abstract/15/3/189>

Hukkanen J, Jacob P, 3rd, and Benowitz NL. Metabolism and disposition kinetics of nicotine. *Pharmacological Reviews*, 2005; 57(1):79-115. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15734728>

Pankow JF, Tavakoli AD, Luo W, and Isabelle LM. Percent free base nicotine in the tobacco smoke particulate matter of selected commercial and reference cigarettes. *Chemical Research in Toxicology*, 2003; 16(8):1014-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12924929>

Jarvis M, Boreham R, Primates P, Feyerabend C, and Bryant A. Nicotine yield from machine-smoked cigarettes and nicotine intakes in smokers: Evidence from a representative population

survey 2001. Journal of the National Cancer Institute, 2001; 93(2):134-8. Available from:
<https://pubmed.ncbi.nlm.nih.gov/11208883/>

Djordjevic M, Stellman S, and Zang E. Doses of nicotine and lung carcinogens delivered to cigarette smokers. Journal of National Cancer Institute, 2000; 92(2):106–11. Available from:
<http://jnci.oxfordjournals.org/cgi/content/full/92/2/106>

DeNoble V. The search for a nicotine analogue that retains the CNS effect of nicotine without increasing cardio-vascular output. in Promoting a Future Without Tobacco, 11th World Conference on Tobacco or Health. Chicago. 2000.

Fant RV, Henningfield JE, Nelson RA, and Pickworth WB. Pharmacokinetics and pharmacodynamics of moist snuff in humans. Tobacco Control, 1999; 8(4):387-92. Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/10629244>

Tomar SL and Henningfield JE. Review of the evidence that pH is a determinant of nicotine dosage from oral use of smokeless tobacco. Tobacco Control, 1997; 6(3):219-25. Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/9396107>

12.4.3.2 Aldehydes

Alamil, H, Colsoul, ML, Heutte, N, Van Der Schueren, M, Galanti, L, & Lechevrel, M. (2024). Exocyclic DNA adducts and oxidative stress parameters: useful tools for biomonitoring exposure to aldehydes in smokers. *Biomarkers*, 1-11. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38506499>

Peterson, LA, Stanfill, SB, & Hecht, SS. (2024). An Update on the Formation in Tobacco, Toxicity, and Carcinogenicity of N¹-Nitrososornicotine and 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38437625>

Wendt, CH, Bowler, RP, Demorest, C, Hastie, A, Labaki, WW, Chen, M et al. (2023). Levels of Urinary Mercapturic Acids of Acrolein, Methacrolein, Crotonaldehyde, and Methyl Vinyl Ketone in Relationship to Chronic Obstructive Pulmonary Disease in Cigarette Smokers of the Subpopulations and Intermediate Outcome Measures in COPD Study (SPIROMICS). *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37725788>

Pavithra, D, Dineshkumar, T, & Rajkumar, K. (2022). Salivary and serum acetaldehyde levels in betel quid chewers, pan masala chewers with or without tobacco and tobacco smokers: A comparative study using head space gas chromatography. *J Oral Maxillofac Pathol*, 26(4), 458-463. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37082040>

Young, M, Feng, C, Cecil, T, & Johnson, TL. (2023). Carbonyl Yields in Cigars under Three Smoking Regimens Using a Linear Smoking Machine. *Chem Res Toxicol*, 36(1), 94-103. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36602460>

Tulen, CBM, Duistermaat, E, Cremers, J, Klerx, WNM, Fokkens, PHB, Weibolt, N et al. (2022). Smoking-Associated Exposure of Human Primary Bronchial Epithelial Cells to Aldehydes: Impact on Molecular Mechanisms Controlling Mitochondrial Content and Function. *Cells*, 11(21). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36359877>

Chen, HC, Cheng, SW, Chen, NY, & Wu, DC. (2022). Characterization and Quantification of Acrolein-Induced Modifications in Hemoglobin by Mass Spectrometry horizontal line Effect of Cigarette Smoking. *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36367988>

Peterson, LA, Seabloom, D, Smith, WE, Vevang, KR, Seelig, DM, Zhang, L, & Wiedmann, TS. (2022). Acrolein Increases the Pulmonary Tumorigenic Activity of the Tobacco-Specific Nitrosamine 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36149460>

Qin Y, Wang B, Liu S, Pan L, Chen M, et al. Robust, comprehensive, sensitive analysis of flavour additives with carboxyl and hydroxyl groups in cigarette smoke combining silylation and gas chromatography-tandem mass spectrometry with an improved backflushing system. *Journal of Chromatography A*, 2022; 1675:463171. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35623195>

Horinouchi T, Mazaki Y, Terada K, and Miwa S. Cigarette smoke extract and its cytotoxic factor acrolein inhibit nitric oxide production in human vascular endothelial cells. *Biol Pharm Bull*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32879145>

World Health Organization. Report on the scientific basis of tobacco product regulation: Seventh report of a WHO study group. Geneva: WHO, 2019. Available from: <https://www.who.int/publications/i/item/who-study-group-on-tobacco-product-regulation-report-on-the-scientific-basis-of-tobacco-product-regulation-seventh-report-of-a-who-study-group>.

Salman R, Talih S, El-Hage R, Haddad C, Karaoghlanian N, et al. Free-base and total nicotine, reactive oxygen species, and carbonyl emissions from iqos, a heated tobacco product. *Nicotine & Tobacco Research*, 2019; 21(9):1285-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30476301>

Jablonski JJ, Maines JH, Cheetham AG, and Gillman IG. Comparative levels of carbonyl delivery between mass-market cigars and cigarettes. *Regulatory Toxicology and Pharmacology*, 2019; 108:104453. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31473262>

Pauwels C, Klerx WNM, Pennings JLA, Boots AW, van Schooten FJ, et al. Cigarette filter ventilation and smoking protocol influence aldehyde smoke yields. *Chemical Research in Toxicology*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29727173>

Bagchi P, Geldner N, deCastro BR, De Jesus VR, Park SK, et al. Crotonaldehyde exposure in U.S. Tobacco smokers and nonsmokers: Nhanes 2005-2006 and 2011-2012. *Environmental research*, 2018; 163:1-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29407484>

Zhao W, Zhang Q, Lu B, Sun S, Zhang S, et al. Rapid determination of six low molecular carbonyl compounds in Tobacco smoke by the APCI-MS/MS coupled to data mining. *Journal of Analytical Methods in Chemistry*, 2017; 2017:8260860. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28512594>

Zhang S, Chen H, Wang A, Liu Y, Hou H, et al. Assessment of genotoxicity of four volatile pollutants from cigarette smoke based on the in vitro gammaH2ax assay using high content screening. *Environmental Toxicology and Pharmacology*, 2017; 55:30-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28818740>

Reilly SM, Goel R, Trushin N, Elias RJ, Foulds J, et al. Brand variation in oxidant production in mainstream cigarette smoke: Carbonyls and free radicals. *Food and Chemical Toxicology*, 2017; 106(Pt A):147-54. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28528972>

Reilly SM, Goel R, Bitzer Z, Elias RJ, Foulds J, et al. Effects of topography-related puff parameters on carbonyl delivery in mainstream cigarette smoke. *Chemical Research in Toxicology*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28648066>

Brancato A, Lavanco G, Cavallaro A, Plescia F, and Cannizzaro C. Acetaldehyde, motivation and stress: Behavioral evidence of an addictive menage a trois. *Frontiers in Behavioral Neuroscience*, 2017; 11:23. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28232795>

Horiyama S, Kunitomo M, Yoshikawa N, and Nakamura K. Mass spectrometric approaches to the identification of potential ingredients in cigarette smoke causing cytotoxicity. *Biol Pharm Bull*, 2016; 39(6):903-8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27251491>

Horiyama S, Hatai M, Takahashi Y, Date S, Masujima T, et al. Intracellular metabolism of alpha,beta-unsaturated carbonyl compounds, acrolein, crotonaldehyde and methyl vinyl ketone, active toxicants in cigarette smoke: Participation of glutathione conjugation ability and aldehyde-ketone sensitive reductase activity. *Chem Pharm Bull (Tokyo)*, 2016; 64(6):585-93. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27250793>

Horinouchi T, Higashi T, Mazaki Y, and Miwa S. Carbonyl compounds in the gas phase of cigarette mainstream smoke and their pharmacological properties. *Biol Pharm Bull*, 2016; 39(6):909-14. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27251492>

Ding YS, Yan X, Wong J, Chan M, and Watson CH. In situ derivatization and quantification of seven carbonyls in cigarette mainstream smoke. *Chemical Research in Toxicology*, 2016; 29(1):125-31. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26700249>

Corley RA, Kabilan S, Kuprat AP, Carson JP, Jacob RE, et al. Comparative risks of aldehyde constituents in cigarette smoke using transient computational fluid dynamics/physiologically based pharmacokinetic models of the rat and human respiratory tracts. *Toxicol Sci*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25858911>

Alwis KU, deCastro BR, Morrow JC, and Blount BC. Acrolein exposure in U.S. Tobacco smokers and non-Tobacco users: Nhanes 2005-2006. *Environmental Health Perspectives*, 2015; 123(12):1302-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26024353>

Pang X and Lewis AC. Carbonyl compounds in gas and particle phases of mainstream cigarette smoke. *Sci Total Environ*, 2011; 409(23):5000-9. Available from: <https://pubmed.ncbi.nlm.nih.gov/21925713/>

Wang M, Cheng G, Balbo S, Carmella S, Villalta P, et al. Clear differences in levels of a formaldehyde-DNA adduct in leukocytes of smokers and nonsmokers. *Cancer Research*, 2009; 69(18):7170-4. Available from: <http://cancerres.aacrjournals.org/cgi/content/full/69/18/7170>

Stevens JF and Maier CS. Acrolein: Sources, metabolism, and biomolecular interactions relevant to human health and disease. *Molecular Nutrition & Food Research*, 2008; 52(1):7-25. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18203133>

Lee EJ, Kim JR, Choi DR, and Ahn YJ. Toxicity of cassia and cinnamon oil compounds and cinnamaldehyde-related compounds to *Sitophilus oryzae* (Coleoptera: Curculionidae). *Journal of Economic Entomology*, 2008; 101(6):1960-6. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/19133480>

Carmella SG, Chen M, Zhang Y, Zhang S, Hatsukami DK, et al. Quantitation of acrolein-derived (3-hydroxypropyl)mercapturic acid in human urine by liquid chromatography-atmospheric pressure chemical ionization tandem mass spectrometry: Effects of cigarette smoking. *Chemical Research in Toxicology*, 2007; 20(7):986-90. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/17559234>

Seeman JI, Laffoon SW, and Kassman AJ. Evaluation of relationships between mainstream smoke acetaldehyde and "tar" and carbon monoxide yields in tobacco smoke and reducing sugars in tobacco blends of U.S. Commercial cigarettes. *Inhalation Toxicology*, 2003; 15(4):373-95. Available from: <https://pubmed.ncbi.nlm.nih.gov/12635005/>

Powrie W, Wu, C.H., Molund, P. Browning reaction systems as sources of mutagens and antimutagens. *Environmental Health Perspectives*, 1986; 67:47-54.

12.4.3.3 Aromatic amines

Ji, H, Jin, Z, Fenton, L, & Slone, S. (2023). Evaluation of Six Aromatic Amines in the Mainstream Smoke of Commercial Cigars. *Chem Res Toxicol*. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/38014781>

Konorev, D, Bellamri, M, Wu, CF, Wu, MT, & Turesky, RJ. (2023). High-Field Asymmetric Waveform Ion Mobility Spectrometry Analysis of Carcinogenic Aromatic Amines in Tobacco Smoke with an Orbitrap Tribrid Mass Spectrometer. *Chem Res Toxicol*. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/37462928>

Scherer, G, Riedel, K, Pluym, N, & Scherer, M. (2022). Assessment of the Exposure to Aromatic Amines in Users of Various Tobacco/Nicotine Products. *ACS Omega*, 7(45), 41775-41782. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36406517>

Ji H and Jin Z. Analysis of six aromatic amines in the mainstream smoke of tobacco products. *Analytical and Bioanalytical Chemistry*, 2022. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/35410388>

Ai J, Hassink M, Taylor KM, Deycard VN, Hearn B, et al. Hydrogen cyanide and aromatic amine yields in the mainstream smoke of 60 little cigars. *Chemical Research in Toxicology*, 2022. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/35612471>

Zhang J, Bai R, Zhou Z, Liu X, and Zhou J. Simultaneous analysis of nine aromatic amines in mainstream cigarette smoke using online solid-phase extraction combined with liquid chromatography-tandem mass spectrometry. *Analytical and Bioanalytical Chemistry*, 2017; 409(11):2993-3005. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28265711>

Deng H, Yang F, Li Z, Bian Z, Fan Z, et al. Rapid determination of 9 aromatic amines in mainstream cigarette smoke by modified dispersive liquid-liquid microextraction and ultraperformance convergence chromatography tandem mass spectrometry. *Journal of Chromatography A*, 2017; 1507:37-44. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28583391>

US Department of Health and Human Services, The health consequences of smoking: A report of the Surgeon General. Atlanta, Georgia: US Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health 2004. Available from:

http://www.cdc.gov/tobacco/data_statistics/sgr/index.htm.

Talaska G. Aromatic amines and human urinary bladder cancer: Exposure sources and epidemiology. *Journal of environmental science and health. Part C, Environmental carcinogenesis & ecotoxicology reviews*, 2003; 21(1):29-43. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12826031>

Grimmer G, Naujack KW, and Dettbarn G. Gaschromatographic determination of polycyclic aromatic hydrocarbons, aza-arenes, aromatic amines in the particle and vapor phase of mainstream and sidestream smoke of cigarettes. *Toxicology letters*, 1987; 35(1):117-24. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/3810672>

12.4.3.4 Hydrocarbons

Bhandari, D, Zhu, Y, Zhang, C, Zhu, W, Alexandridis, A, Etemadi, A et al. (2023). Smoke Exposure Associated with Higher Urinary Benzene Biomarker Muconic Acid (MUCA) in Golestan Cohort Study Participants. *Biomarkers*, 1-9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37878492>

Wu, G, Gong, S, He, Y, & Liu, D. (2023). Smoking is associated with elevated blood level of volatile organic compounds: a population-based analysis of NHANES 2017-2018. *Arch Public Health*, 81(1), 55. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37055810>

Yang, Y, Wang, C, Zhang, H, Qian, J, Yang, S, Liao, H et al. (2023). Preparation of Functionalized Zr-Based MOFs and MOFs/GO for Efficient Removal of 1,3-Butadiene from Cigarette Smoke. *Materials (Basel)*, 16(2). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36676418>

Fiebelkorn S and Meredith C. Estimation of the leukemia risk in human populations exposed to benzene from tobacco smoke using epidemiological data. *Risk Analysis*, 2018; 38(7):1490-501. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29266361>

International Agency for Research on Cancer Working Group on the Evaluation of Carcinogenic Risks to Humans. Tobacco smoke and involuntary smoking. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, Volume 83. Lyon: International Agency for Research on Cancer 2004. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Tobacco-Smoke-And-Involuntary-Smoking-2004>.

U.S. Environmental Protection Agency. Benzene; casrn 71-43-2. Integrated Risk Information System (IRIS) U.S. Environmental Protection Agency Chemical Assessment Summary: EPA, 2003. Available from: https://iris.epa.gov/static/pdfs/0276_summary.pdf.

Fowles J and Dybing E. Application of toxicological risk assessment principles to the chemical constituents of cigarette smoke. *Tobacco Control*, 2003; 12(4):424-30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/14660781>

Korte JE, Hertz-Picciotto I, Schulz MR, Ball LM, and Duell EJ. The contribution of benzene to smoking-induced leukemia. *Environmental Health Perspectives*, 2000; 108(4):333-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/10753092>

National Toxicology Program. Ntp toxicology and carcinogenesis studies of isoprene (cas no. 78-79-5) in f344/n rats (inhalation studies). National Toxicology Program Technical Report Series, 1999; 486:1-176. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12571689>

Brunnemann KD, Kagan MR, Cox JE, and Hoffmann D. Determination of benzene, toluene and 1,3-butadiene in cigarette smoke by GC-MDS. *International Journal of Experimental Pathology*, 1989; 37(1-4):108-13. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/2637141>

Grimmer G, Naujack KW, and Dettbarn G. Gaschromatographic determination of polycyclic aromatic hydrocarbons, aza-arenes, aromatic amines in the particle and vapor phase of mainstream and sidestream smoke of cigarettes. *Toxicology letters*, 1987; 35(1):117-24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/3810672>

12.4.3.5 Heterocyclic aromatic amines

Zhang L, Wang L, Li Y, Xia Y, Chang CM, et al. Evaluation of tobacco smoke and diet as sources of exposure to two heterocyclic aromatic amines for the U.S. Population: Nhanes 2013-2014. *Cancer Epidemiology, Biomarkers & Prevention*, 2020; 29(1):103-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31575556>

Roemer E, Meisgen T, Diekmann J, Conroy L, and Stabbert R. Heterocyclic aromatic amines and their contribution to the bacterial mutagenicity of the particulate phase of cigarette smoke. *Toxicology letters*, 2016; 243:40-7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26724587>

Turesky RJ. Heterocyclic aromatic amine metabolism, DNA adduct formation, mutagenesis, and carcinogenesis. *Drug Metabolism Reviews*, 2002; 34(3):625-50. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12214671>

Sugimura T. Overview of carcinogenic heterocyclic amines. *Mutation Research*, 1997; 376(1-2):211-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/9202758>

Nagao M and Sugimura T. Carcinogenic factors in food with relevance to colon cancer development. *Mutation Research*, 1993; 290(1):43-51. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/7694098>

12.4.3.6 Polycyclic aromatic hydrocarbons (PAH)

Tang, H, Yuan, K, Chu, F, Zhang, X, Li, Q, Chen, Q et al. (2024). On-Line Analysis of Cigarette Smoke Based on Microwave Plasma Torch Mass Spectrometry. *ChemistryOpen*, e202400013. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38873881>

Tao, L, Chiarelli, MP, Pavlova, S, Kolokythas, A, Schwartz, J, DeFrancesco, J et al. (2024). Enrichment of polycyclic aromatic hydrocarbon metabolizing microorganisms on the oral mucosa of tobacco users. *PeerJ*, 12, e16626. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38188172>

Gao, B, Ma, S, Zhao, W, Tian, M, Huang, Y, Chen, L et al. (2023). Calculated cancer risks for polycyclic aromatic hydrocarbon mixtures in mainstream smoke of cigarettes sold in China. *Regul Toxicol Pharmacol*, 142, 105427. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37290570>

Edwards, SH, Hassink, M, Taylor, KM, & Vu, AT. (2023). Variation of Benzo[a]pyrene, NNN, and NNK Levels in 16 Commercial Smokeless Tobacco Products. *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36638203>

Nsonwu-Anyanwu, AC, Egom, OU, Eworo, RE, Nsonwu, MC, Aniekpon, UF, Ekpo, DO, & Opara Usoro, CA. (2022). Risk of Pulmonary-Reproductive Dysfunctions, Inflammation and Oxidative DNA Damage in Exposure to Polycyclic Aromatic Hydrocarbon in Cigarette Smokers. *Med J Islam Repub Iran*, 36, 108. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36447550>

Abadi, DRV, Tahmasbizadeh, M, Arfaenia, H, Masjedi, MR, Ramavandi, B, & Poureshgh, Y. (2022). Biomonitoring of unmetabolized polycyclic aromatic hydrocarbons (PAHs) in urine of waterpipe/cigarette cafe workers. *Environ Sci Pollut Res Int*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36306072>

Chang, J, Wang, Q, Dong, X, Luo, T, Liu, Z, & Xu, D. (2022). The influencing factors of health hazards of benzo[a]pyrene in cigarette mainstream smoke: The example of one brand in Beijing. *Tob Induc Dis*, 20, 80. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36212736>

Wang SQ, Wang WJ, Wu CC, Bao LJ, Yu Y, et al. Low tar level does not reduce human exposure to polycyclic aromatic hydrocarbons in environmental Tobacco smoke. *Environmental Science & Technology*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31859494>

Jeffery J, Carradus M, Songin K, Pettit M, Pettit K, et al. Optimized method for determination of 16 FDA polycyclic aromatic hydrocarbons (pahs) in mainstream cigarette smoke by gas chromatography-mass spectrometry. *Chemistry Central Journal*, 2018; 12(1):27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29536204>

Zhang Y, Zou HY, Shi P, Yang Q, Tang LJ, et al. Determination of benzo[a]pyrene in cigarette mainstream smoke by using mid-infrared spectroscopy associated with a novel chemometric algorithm. *Analytica Chimica Acta*, 2016; 902:43-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26703252>

Rodgman A and Perfetti T. The composition of cigarette smoke: A catalogue of the polycyclic aromatic hydrocarbons. *Eiträge zur Tabakforschung International*, 2016; 22(1):13-69. Available from: <https://cyberleninka.org/article/n/650361/viewer>

Paschke M, Hutzler C, Henkler F, and Luch A. Oxidative and inert pyrolysis on-line coupled to gas chromatography with mass spectrometric detection: On the pyrolysis products of tobacco additives. *Int J Hyg Environ Health*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27622657>

Ohkubo T, Inaba Y, Hara Y, Uchiyama S, and Kunugita N. Determination of polycyclic aromatic hydrocarbons and mutagenicity of mainstream smoke and heavy metals in Tobacco filler of cigarettes of a brand in Japan and cigarettes of the same brand imported privately from other asian countries. *Nihon Eiseigaku Zasshi*, 2016; 71(1):84-90. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26832621>

Luo YB, Chen XJ, Zhang HF, Jiang XY, Li X, et al. Simultaneous determination of polycyclic aromatic hydrocarbons and tobacco-specific n-nitrosamines in mainstream cigarette smoke using in-pipette-tip solid-phase extraction and on-line gel permeation chromatography-gas chromatography-tandem mass spectrometry. *Journal of Chromatography A*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27435688>

Zhang X, Hou H, Chen H, Liu Y, Wang A, et al. Quantification of 16 polycyclic aromatic hydrocarbons in cigarette smoke condensate using stable isotope dilution liquid chromatography with atmospheric-pressure photoionization tandem mass spectrometry. *Journal of Separation Science*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26377484>

Vu AT, Taylor KM, Holman MR, Ding YS, Hearn B, et al. Polycyclic aromatic hydrocarbons in the mainstream smoke of popular U.S. Cigarettes. *Chemical Research in Toxicology*, 2015; 28(8):1616-26. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26158771>

Tobacco Laboratory Network. Determination of benzo[a]pyrene in mainstream cigarette smoke under ISO and intense smoking conditions. WHO TobLabNet Official Method, SOP 05. Geneva: WHO, 2015. Available from: <https://apps.who.int/iris/rest/bitstreams/783899/retrieve>.

Gao B, Du X, Wang X, Tang J, Ding X, et al. Parent, alkylated, and sulfur/oxygen-containing polycyclic aromatic hydrocarbons in mainstream smoke from 13 brands of chinese cigarettes. *Environmental Science & Technology*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26119395>

Lee HL, Hsieh DP, and Li LA. Polycyclic aromatic hydrocarbons in cigarette sidestream smoke particulates from a Taiwanese brand and their carcinogenic relevance. *Chemosphere*, 2011; 82(3):477-82. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20947129>

Zhong Y, Carmella S, Upadhyaya P, Hochalter J, Rauch D, et al. Immediate consequences of cigarette smoking: Rapid formation of polycyclic aromatic hydrocarbon diol epoxides. *Chemical Research in Toxicology*, 2010; [Epub ahead of print]. Available from: <http://pubs.acs.org/doi/full/10.1021/tx100345x>

International Agency for Research on Cancer. IARC monographs on the evaluation of carcinogenic risks to humans: Some non-heterocyclic polycyclic aromatic hydrocarbons and some related exposures. *Monographs on the Evaluation of Carcinogenic Risks to Human*, 92 Lyon, France: IARC, 2010. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Some-Non-heterocyclic-Polycyclic-Aromatic-Hydrocarbons-And-Some-Related-Exposures-2010>.

Choi H, Harrison, R., Komulainen, H., Delgado-Saborit, J.M. Polycyclic aromatic hydrocarbons. WHO Guidelines for Indoor Air Quality: Selected Pollutants., Geneva: World Health Organization., 2010.

Watanabe K, Djordjevic M, Stellman S, Toccalino P, Austin D, et al. Incremental lifetime cancer risks computed for benzo[a]pyrene and two tobacco-specific n-nitrosamines in mainstream cigarette smoke compared with lung cancer risks derived from epidemiologic data. *Regulatory Toxicology and Pharmacology* 2009; 57(1):24–30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20025920>

Armstrong B and Gibbs G. Exposure-response relationship between lung cancer and polycyclic aromatic hydrocarbons (pahs). *Occupational and Environmental Medicine*, 2009; 66(11):740–6. Available from: <http://oem.bmj.com/cgi/content/full/66/11/740>

Monzer B, Sepetdjian E, Saliba N, and Shihadeh A. Charcoal emissions as a source of co and carcinogenic pah in mainstream narghile waterpipe smoke. *Food and Chemical Toxicology*, 2008; 46(9):2991-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18573302>

Ding YS, Zhang L, Jain RB, Jain N, Wang RY, et al. Levels of tobacco-specific nitrosamines and polycyclic aromatic hydrocarbons in mainstream smoke from different tobacco varieties. *Cancer Epidemiology, Biomarkers & Prevention*, 2008; 17(12):3366-71. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19064552>

Akpan V, Huang S, Lodovici M, and Dolara P. High levels of carcinogenic polycyclic aromatic hydrocarbons (pah) in 20 brands of chinese cigarettes. *Journal of Applied Toxicology*, 2006; 26(6):480-3. Available from: <https://pubmed.ncbi.nlm.nih.gov/17080399/>

Ding YS, Trommel JS, Yan XJ, Ashley D, and Watson CH. Determination of 14 polycyclic aromatic hydrocarbons in mainstream smoke from domestic cigarettes. *Environmental Science & Technology*, 2005; 39(2):471-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15707046>

Brunnemann KD, Kagan MR, Cox JE, and Hoffmann D. Analysis of 1,3-butadiene and other selected gas-phase components in cigarette mainstream and sidestream smoke by gas chromatography-mass selective detection. *Carcinogenesis*, 1990; 11(10):1863-8. Available from: <https://pubmed.ncbi.nlm.nih.gov/2208599/>

Vu-Duc T and Cong-Khanh H. Sidestream tobacco smoke constituents in indoor air modelled in an experimental chamber: Polycyclic aromatic hydrocarbons. *Environment International*, 1989; 15:57 - 64. Available from: <https://www.sciencedirect.com/science/article/pii/016041208990010X>

Grimmer G, Naujack KW, and Dettbarn G. Gaschromatographic determination of polycyclic aromatic hydrocarbons, aza-arenes, aromatic amines in the particle and vapor phase of mainstream and sidestream smoke of cigarettes. *Toxicology letters*, 1987; 35(1):117-24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/3810672>

12.4.3.7 N-nitrosamines

Peterson, LA, Stanfill, SB, & Hecht, SS. (2024). An Update on the Formation in Tobacco, Toxicity, and Carcinogenicity of N'-Nitrosoornicotine and 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38437625>

Ai, J, Hassink, M, Taylor, KM, Kuklenyik, P, Valentin-Blasini, L, & Watson, C. (2024). Tobacco-Specific Nitrosamines in Current Commercial Large Cigars, Cigarillos, and Little Cigars. *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38235526>

Bitzer, Z, Mocniak, LE, Trushin, N, Smith, M, & Richie, JP. (2023). Influence of Tobacco Variety and Curing on Free Radical Production in Cigarette Smoke. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36967618>

Edwards, SH, Hassink, M, Taylor, KM, & Vu, AT. (2023). Variation of Benzo[a]pyrene, NNN, and NNK Levels in 16 Commercial Smokeless Tobacco Products. *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36638203>

Thasweer, AM, Renuka Devi, P, & Thirunavukkarasu, V. (2022). Molecular docking and dynamic simulation studies of alpha4beta2 and alpha7 nicotinic acetylcholine receptors with tobacco smoke

constituents nicotine, NNK and NNN. *J Biomol Struct Dyn*, 1-10. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36270967>

Peterson, LA, Seabloom, D, Smith, WE, Vevang, KR, Seelig, DM, Zhang, L, & Wiedmann, TS. (2022). Acrolein Increases the Pulmonary Tumorigenic Activity of the Tobacco-Specific Nitrosamine 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36149460>

Zhang J, Liu X, Shi B, Yang Z, Luo Y, et al. Investigation of exposure biomarkers in human plasma following differing levels of tobacco-specific n-nitrosamines and nicotine in cigarette smoke. *Environmental research*, 2022; 214(Pt 2):113811. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35835167>

Tayyarah R. Comments on “Tobacco-specific nitrosamines in the Tobacco and mainstream smoke of commercial little cigars”. *Chemical Research in Toxicology*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35738011>

Sarker AH and Hang B. Tobacco-specific nitrosamine 1-(n-methyl-n-nitrosamino)-1-(3-pyridinyl)-4-butanal (nna) causes DNA damage and impaired replication/transcription in human lung cells. *PLoS One*, 2022; 17(5):e0267839. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35576221>

Lee JW, Yang W, Kim YS, Kim Y, Yoo HS, et al. Exposure to secondhand smoke and a Tobacco-specific carcinogen in non-smokers. *Korean J Fam Med*, 2022; 43(2):117-24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35320897>

Hecht SS, Gupta PC, Sturla SJ, and Wang Y. 50 years of research on Tobacco-specific nitrosamines: A virtual collection of emerging knowledge of chemical toxicology of Tobacco and nicotine delivery systems and call for contributions to a landmark special issue. *Chemical Research in Toxicology*, 2022; 35(6):899-900. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35635503>

Edwards SH, Hassink MD, Taylor KM, Watson CH, Kuklennyik P, et al. Response to ‘comments on “Tobacco-specific nitrosamines in the Tobacco and mainstream smoke of commercial little cigars”’ letter to editor. *Chemical Research in Toxicology*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35738012>

Cheng CW, Kou HS, Wu SM, and Wang CC. A chemometric experimental design with three-step stacking capillary electrophoresis for analysis of five tobacco-specific nitrosamines in cigarette products. *Journal of Chromatography A*, 2022; 1677:463283. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35810639>

Xia B, Blount BC, Guillot T, Brosius C, Li Y, et al. Tobacco-specific nitrosamines (NNAL, nnn, nat, and nab) exposures in the US population assessment of Tobacco and health (path) study wave 1 (2013-2014). *Nicotine & Tobacco Research*, 2021; 23(3):573-83. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32716026>

Guo J, Chen H, Upadhyaya P, Zhao Y, Turesky RJ, et al. Mass spectrometric quantitation of apurinic/aprimidinic sites in tissue DNA of rats exposed to tobacco-specific nitrosamines and in lung and leukocyte DNA of cigarette smokers and non-smokers. *Chemical Research in Toxicology*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32833447>

Ishizaki A and Kataoka H. A sensitive method for the determination of tobacco-specific nitrosamines in mainstream and sidestream smokes of combustion cigarettes and heated tobacco products by online in-tube solid-phase microextraction coupled with liquid chromatography-tandem mass spectrometry. *Analytica Chimica Acta*, 2019; 1075:98-105. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31196428>

Zhang S, Chen H, Wang A, Liu Y, Hou H, et al. Genotoxicity analysis of five particle matter toxicants from cigarette smoke based on gammaH2ax assay combined with hill/two-component model. *Environ Toxicol Pharmacol*, 2018; 58:131-40. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29329021>

Tobacco laboratory Network. Determination of tobacco-specific nitrosamines in mainstream cigarette smoke under ISO and intense smoking conditions., SOP 08. Geneva: WHO, 2018. Available from: <https://apps.who.int/iris/rest/bitstreams/1156182/retrieve>.

Leigh NJ, Palumbo MN, Marino AM, O'Connor RJ, and Goniewicz ML. Tobacco-specific nitrosamines (tsna) in heated tobacco product iqos. *Tobacco Control*, 2018; 27(Suppl 1):s37-s8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30242043>

Konstantinou E, Fotopoulou F, Drosos A, Dimakopoulou N, Zagoriti Z, et al. Tobacco-specific nitrosamines: A literature review. *Food and Chemical Toxicology*, 2018; 118:198-203. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29751076>

Gushgari AJ and Halden RU. Critical review of major sources of human exposure to n-nitrosamines. *Chemosphere*, 2018; 210:1124-36. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30208538>

Czoli CD and Hammond D. Carcinogen exposure among canadian tobacco users: Changes in NNK exposure from 2007-09 through 2012-13. *Cancer Epidemiology, Biomarkers & Prevention*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29358222>

Khariwala SS, Ma B, Ruszczak C, Carmella SG, Lindgren B, et al. High level of tobacco carcinogen-derived DNA damage in oral cells is an independent predictor of oral/head and neck cancer risk in smokers. *Cancer Prevention Research (Philadelphia, Pa.)*, 2017; 10(9):507-13. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28679497>

Zhang J, Bai R, Yi X, Yang Z, Liu X, et al. Fully automated analysis of four tobacco-specific n-nitrosamines in mainstream cigarette smoke using two-dimensional online solid phase extraction combined with liquid chromatography-tandem mass spectrometry. *Talanta*, 2016; 146:216-24. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26695255>

Yalcin E and de la Monte S. Tobacco nitrosamines as culprits in disease: Mechanisms reviewed. *Journal of Physiology and Biochemistry*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26767836>

Luo YB, Chen XJ, Zhang HF, Jiang XY, Li X, et al. Simultaneous determination of polycyclic aromatic hydrocarbons and tobacco-specific n-nitrosamines in mainstream cigarette smoke using in-pipette-tip solid-phase extraction and on-line gel permeation chromatography-gas chromatography-tandem mass spectrometry. *Journal of Chromatography A*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27435688>

Gunduz I, Kondylis A, Jaccard G, Renaud JM, Hofer R, et al. Tobacco-specific n-nitrosamines nnn and NNK levels in cigarette brands between 2000 and 2014. *Regulatory Toxicology and Pharmacology*, 2016; 76:113-20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26806560>

Xiang Z, Cai K, Zhou S, Geng Z, and Pan W. Analysis of nitrogenous organic compounds from mainstream cigarette smoke using low-temperature solvent extraction followed by comprehensive two-dimensional gas chromatography with high-resolution time-of-flight mass spectrometry. *Journal of Separation Science*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25962488>

Hyodo T, Minagawa K, and Mikita A. Five-year yield variation in n-nitrosornicotine and (4-methylnitrosoamino)-1-(3-pyridyl)-1-butanone from the smoke of commercial cigarette brands on the Japanese market. *Regulatory Toxicology and Pharmacology*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26382610>

Tobacco Laboratory Network. Standard operating procedure for determination of tobacco-specific nitrosamines in mainstream cigarette smoke under ISO and intense smoking conditions. WHO TobLabNet Official Method, SOP 03. Geneva: WHO, 2014. Available from: https://apps.who.int/iris/bitstream/handle/10665/136000/9789241506663_eng.pdf;sequence=1.

Shah K and Karnes H. A review of the analysis of tobacco-specific nitrosamines in biological matrices. *Critical Reviews in Toxicology*, 2010; 40(4):305–27. Available from: <http://informahealthcare.com/doi/pdf/10.3109/10408440903394435>

Watanabe K, Djordjevic M, Stellman S, Toccalino P, Austin D, et al. Incremental lifetime cancer risks computed for benzo[a]pyrene and two tobacco-specific n-nitrosamines in mainstream cigarette smoke compared with lung cancer risks derived from epidemiologic data. *Regulatory Toxicology and Pharmacology* 2009; 57(1):24–30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20025920>

Polzin G, Wu W, Yan X, McCraw J, Abdul-Salaam S, et al. Estimating smokers' mouth-level exposure to select mainstream smoke constituents from discarded cigarette filter butts. *Nicotine & Tobacco Research*, 2009; 11(7):868–74. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19541951

Nishioka T, Guo J, Yamamoto D, Chen L, Huppi P, et al. Nicotine, through upregulating pro-survival signaling, cooperates with NNK to promote transformation. *Journal of Cellular Biochemistry*, 2009; 109(1):152–61. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/jcb.22392/full>

Kavvadias D, Scherer G, Cheung F, Errington G, Shepperd J, et al. Determination of tobacco-specific n-nitrosamines in urine of smokers and non-smokers. *Biomarkers : Biochemical Indicators of Exposure, Response, and Susceptibility to Chemicals*, 2009; 14(8). Available from: <http://informahealthcare.com/doi/full/10.3109/13547500903242883>

Rickert W, Joza P, Sharifi M, Wu J, and Lauterbach J. Reductions in the tobacco specific nitrosamine (tsna) content of tobaccos taken from commercial Canadian cigarettes and corresponding reductions in tsna deliveries in mainstream smoke from such cigarettes. *Regulatory Toxicology And Pharmacology*, 2008; 51(3):306–10. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18508168>

- Li M-Y, Yip J, Hsin M, Mok T, Wu Y, et al. Ho-1 plays a central role in NNK-mediated lung carcinogenesis. *The European Respiratory Journal*, 2008; 32(4):911–23. Available from: <http://erj.ersjournals.com/content/32/4/911.long>
- Hertsgaard L, Hanson K, Hecht S, Lindgren B, Luo X, et al. Exposure to a tobacco-specific lung carcinogen in adolescent versus adult smokers. *Cancer Epidemiology, Biomarkers & Prevention* 2008; 17(12):3337–43. Available from: <http://cebp.aacrjournals.org/cgi/content/full/17/12/3337>
- Gartner CE and Hall WD. Should Australia lift its ban on low nitrosamine smokeless tobacco products? *Medical Journal of Australia*, 2008; 188(1):44-6. Available from: http://www.mja.com.au/public/issues/188_01_070108/gar10502_fm.html
- Ding YS, Zhang L, Jain RB, Jain N, Wang RY, et al. Levels of tobacco-specific nitrosamines and polycyclic aromatic hydrocarbons in mainstream smoke from different tobacco varieties. *Cancer Epidemiology, Biomarkers & Prevention*, 2008; 17(12):3366-71. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19064552>
- International Agency for Research on Cancer. IARC monographs on the evaluation of carcinogenic risks to humans: Smokeless tobacco and some tobacco-specific n-nitrosamines. 89 Lyon, France: IARC, 2007. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Smokeless-Tobacco-And-Some-Tobacco-specific-Em-N-Em--Nitrosamines-2007>.
- Gray N and Boyle P. The case of the disappearing nitrosamines: A potentially global phenomenon. *Tobacco Control*, 2004; 13(1):13-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/14985588>
- Breland A, Acosta M, and Eissenberg T. Tobacco specific nitrosamines and potential reduced exposure products for smokers: A preliminary evaluation of advancetm. *Tobacco Control*, 2003; 12:317-21. Available from: <http://tobaccocontrol.bmj.com/cgi/content/abstract/12/3/317>
- Ashley D, Johnson DR, McCraw JM, Richter P, Pirkle JL, et al. Tobacco-specific nitrosamines in U.S. Brand and non-U.S. Brand cigarettes. *Nicotine and Tobacco Research*, 2003; 5(3):323-32. Available from: <https://pubmed.ncbi.nlm.nih.gov/12791527/>
- Gray N, Zaridze D, Robertson C, Krivosheeva L, Sigacheva N, et al. Variation within global cigarette brands in tar, nicotine, and certain nitrosamines: Analytic study. *Tobacco Control*, 2000; 9(3):351. Available from: <http://tobaccocontrol.bmj.com/cgi/content/full/9/3/351>
- Klus H, Begutter H, Scherer G, Tricker AR, and Adlkofer F. Tobacco-specific and volatile n-nitrosamines in environmental Tobacco smoke of offices. *Indoor Environment*, 1992; 1(6):348 - 50. Available from: <https://journals.sagepub.com/doi/abs/10.1177/1420326X9200100606>
- Brunnemann KD, Cox JE, and Hoffmann D. Analysis of tobacco-specific n-nitrosamines in indoor air. *Carcinogenesis*, 1992; 13(12):2415-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/1473252>
- Fisher S, Spiegelhalder B, and Eisenbarth J. Investigations on the origins of tobacco-specific nitrosamines in mainstream smoke of cigarettes. *Carcinogenesis*, 1990; 11(5):723–30. Available from: <http://carcin.oxfordjournals.org/cgi/reprint/11/5/723>

Ohshima H, Nair J, Bourgade MC, Friesen M, Garren L, et al. Identification and occurrence of two new n-nitrosamino acids in tobacco products: 3-(N-nitroso-N-methylamino)propionic acid and 4-(N-nitroso-N-methylamino)butyric acid. *Cancer Letters*, 1985; 26(2):153-62. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/3978605>

12.4.3.8 Phenols

Song, X, Qian, S, Li, H, Shen, Y, Bian, D, Shan, K, & Shi, J. (2024). Study on the evaluation method of cigarette astringency in the simulated oral environment. *J Texture Stud*, 55(3), e12837. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38702991>

Si, X, Yang, J, Zhang, F, Zhu, R, Liu, C, Jiang, W et al. (2022). Determination and Difference Analysis of Phenolic Compounds in Smokers' Saliva and Mainstream Smoke. *J Anal Methods Chem*, 2022, 6788394. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36213091>

Ren Z, Nie B, Liu T, Yuan F, Feng F, et al. Simultaneous determination of coumarin and its derivatives in Tobacco products by liquid chromatography-tandem mass spectrometry. *Molecules*, 2016; 21(11). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27834935>

Kibet JK, Khachatryan L, and Dellinger B. Phenols from pyrolysis and co-pyrolysis of tobacco biomass components. *Chemosphere*, 2015; 138:259-65. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26091866>

Saha S, Mistri R, and Ray BC. A rapid and selective method for simultaneous determination of six toxic phenolic compounds in mainstream cigarette smoke using single-drop microextraction followed by liquid chromatography-tandem mass spectrometry. *Analytical and Bioanalytical Chemistry*, 2013; 405(28):9265-72. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24057026>

Wu J, Rickert WS, and Masters A. An improved high performance liquid chromatography-fluorescence detection method for the analysis of major phenolic compounds in cigarette smoke and smokeless tobacco products. *Journal of Chromatography A*, 2012; 1264:40-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23062878>

Vaughan C, Stanfill SB, Polzin GM, Ashley DL, and Watson CH. Automated determination of seven phenolic compounds in mainstream tobacco smoke. *Nicotine & Tobacco Research*, 2008; 10(7):1261-8. Available from: <http://www.informaworld.com/smpp/content~content=a794983022~db=all~order=page>

McGregor D. Hydroquinone: An evaluation of the human risks from its carcinogenic and mutagenic properties. *Critical Reviews in Toxicology*, 2007; 37(10):887-914. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18027166>

Leanderson P and Tagesson C. Cigarette smoke-induced DNA-damage: Role of hydroquinone and catechol in the formation of the oxidative DNA-adduct, 8-hydroxydeoxyguanosine. *Chemico-Biological Interactions*, 1990; 75(1):71-81. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/2114224>

12.4.3.9 Other carbon-based (organic) compounds

Yang, F, Cui, H, Wang, C, Wang, Y, Zhu, W, Deng, H et al (2023). Comparison of supercritical fluid chromatography-tandem mass spectrometry and liquid chromatography-tandem mass spectrometry for the stereoselective analysis of chlorfenvinphos and dimethylvinphos in tobacco. *J Sep Sci*, e2300449. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37582637>

Wu, G, Gong, S, He, Y, & Liu, D. (2023). Smoking is associated with elevated blood level of volatile organic compounds: a population-based analysis of NHANES 2017-2018. *Arch Public Health*, 81(1), 55. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37055810>

Picanco, JMA, Limberger, RP, & Apel, MA. (2022). Uncovering cloves: characterization of volatile compounds present in clove cigarettes. *Toxicol Res (Camb)*, 11(6), 987-1002. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36569486>

Kenwood BM, Zhu W, Zhang L, Bhandari D, and Blount BC. Cigarette smoking is associated with acrylamide exposure among the U.S. Population: Nhanes 2011-2016. *Environmental research*, 2022:112774. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35074357>

Junco JG, Chapman GM, Bravo Cardenas R, Watson CH, and Valentin-Blasini L. Quantification of nitromethane in mainstream smoke using gas chromatography and tandem mass spectrometry. *Toxicology Reports*, 2021; 8:405-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33717993>

Espenship M, Silva LK, Smith MM, Capella KM, Reese CM, et al. Nitromethane exposure from Tobacco smoke and diet in the U.S. Population: Nhanes, 2007 - 2012. *Environmental Science & Technology*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30672285>

Marchese S, Polo A, Ariano A, Velotto S, Costantini S, et al. Aflatoxin b1 and m1: Biological properties and their involvement in cancer development. *Toxins (Basel)*, 2018; 10(6). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29794965>

Mojska H, Gielecinska I, and Cendrowski A. Acrylamide content in cigarette mainstream smoke and estimation of exposure to acrylamide from tobacco smoke in Poland. *Ann Agric Environ Med*, 2016; 23(3):456-61. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27660868>

Xiang Z, Cai K, Zhou S, Geng Z, and Pan W. Analysis of nitrogenous organic compounds from mainstream cigarette smoke using low-temperature solvent extraction followed by comprehensive two-dimensional gas chromatography with high-resolution time-of-flight mass spectrometry. *Journal of Separation Science*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25962488>

Wang X, Guo J, Shang J, Ding L, Zhao G, et al. Determination of nitroalkanes in mainstream cigarette smoke by heart-cutting multidimensional gas chromatography system coupled with mass spectrometry detection. *Journal of Chromatography A*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26603996>

Chagovets V, Kononikhin A, Starodubtseva N, Ding J, Chen H, et al. Letter: Comparison of pyridine and pyrazine derivatives distributions in exhaled breath and exhaled breath condensate after smoking. *Eur J Mass Spectrom (Chichester, Eng)*, 2015; 21(6):829-32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26764312>

Novak BJ, Meinardi S, and Blake DR. Methyl chloride and the U.S. Cigarette. *Nicotine & Tobacco Research*, 2008; 10(11):1621–5. Available from:

<http://www.informaworld.com/smpp/content~content=a905081554~db=all~order=page>

International Agency for Research on Cancer. Monographs on the evaluation of carcinogenic risks to humans: Some naturally occurring substances: Food items and constituents, heterocyclic aromatic amines and mycotoxins., 56 Lyon, France: IARC, 1993. Available from:

<https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Some-Naturally-Occurring-Substances-Food-Items-And-Constituents-Heterocyclic-Aromatic-Amines-And-Mycotoxins-1993>.

12.4.3.10 Inorganics

Mocniak, L., Bitzer, Z., Goel, R, Muscat, JE, Foulds, J, Elias, RJ, & Richie, JP. (2024). Free Radicals in Little Cigar Mainstream Smoke and the Potential Influence of Flavoring Chemicals on Free Radical Production. *Chem Res Toxicol*, 37(7), 1121-1128. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/38953874>

Mehrparvar AH, Ghanbari L, Mirmohammadi SJ, Zare Sakhvidi MJ, Vakili M, et al. The effect of water-pipe and cigarette smoking on exhaled nitric oxide. *International Journal of Preventive Medicine*, 2022; 13:79. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35706853>

Ai J, Hassink M, Taylor KM, Deycard VN, Hearn B, et al. Hydrogen cyanide and aromatic amine yields in the mainstream smoke of 60 little cigars. *Chemical Research in Toxicology*, 2022. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/35612471>

Merianos AL, Jandarov RA, Cataletto M, and Mahabee-Gittens EM. Tobacco smoke exposure and fractional exhaled nitric oxide levels among U.S. Adolescents. *Nitric Oxide*, 2021; 117:53-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34688860>

Zaga V, Cattaruzza MS, Martucci P, Pacifici R, Trisolini R, et al. The “polonium in vivo” study: Polonium-210 in bronchial lavages of patients with suspected lung cancer. *Biomedicine*, 2020; 9(1). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33374630>

Sharifi M, Donisa C, and Joza P. A sensitive and quantitative isotope-dilution lc-ms/ms method for analysis of hydrazine in Tobacco smoke. *Journal of Chromatographic Science*, 2019. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31711231>

Laking GR. Human exposure to radioactivity from tobacco smoke: Systematic review. *Nicotine & Tobacco Research*, 2019; 21(9):1172-80. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30060241>

Dorey A, Scheerlinck P, Nguyen H, and Albertson T. Acute and chronic carbon monoxide toxicity from Tobacco smoking. *Mil Med*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31670370>

Inaba Y, Uchiyama S, and Kunugita N. Spectrophotometric determination of ammonia levels in tobacco fillers of and sidestream smoke from different cigarette brands in Japan. *Environmental Health and Preventive Medicine*, 2018; 23(1):15. Available from:

<https://pubmed.ncbi.nlm.nih.gov/29703135/>

Inaba Y, Uchiyama S, and Kunugita N. Spectrophotometric determination of ammonia levels in tobacco fillers of and sidestream smoke from different cigarette brands in Japan. *Environmental Health and Preventive Medicine*, 2018; 23(1):15. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29703135>

Watson CV, Feng J, Valentin-Blasini L, Stanelle R, and Watson CH. Method for the determination of ammonia in mainstream cigarette smoke using ion chromatography. *PLoS One*, 2016; 11(7):e0159126. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27415766>

Wang Y, Liu M, Zhu Y, Cheng K, Da W, et al. Identifying the tobacco related free radicals by upcc-qtof-ms with radical trapping method in mainstream cigarette smoke. *Talanta*, 2016; 160:106-12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27591593>

Tobacco laboratory Network. Determination of nicotine and carbon monoxide in mainstream cigarette smoke under intense smoking conditions. SOP 10. Geneva: WHO, 2016. Available from: <https://www.who.int/publications/i/item/standard-operating-procedure-for-determination-of-nicotine-and-carbon-monoxide-in-mainstream-cigarette-smoke-under-intense-smoking-conditions>.

Watson CV, Valentin-Blasini L, Damian M, and Watson CH. Method for the determination of ammonium in cigarette tobacco using ion chromatography. *Regulatory Toxicology and Pharmacology*, 2015; 72(2):266-70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25934256>

Mahernia S, Amanlou A, Kiaee G, and Amanlou M. Determination of hydrogen cyanide concentration in mainstream smoke of tobacco products by polarography. *J Environ Health Sci Eng*, 2015; 13:57. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26225214>

Gondal MA, Habibullah YB, Oloore LE, and Iqbal MA. Determination of carcinogenic fluorine in cigarettes using pulsed uv laser-induced breakdown spectroscopy. *Appl Opt*, 2015; 54(17):5560-7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26192861>

Zhang ZW, Xu YB, Wang CH, Chen KB, Tong HW, et al. Direct determination of hydrogen cyanide in cigarette mainstream smoke by ion chromatography with pulsed amperometric detection. *Journal of Chromatography A*, 2011; 1218(7):1016-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21238972>

van Amsterdam J, Sleijffers A, van Spiegel P, Blom R, Witte M, et al. Effect of ammonia in cigarette tobacco on nicotine absorption in human smokers. *Food and Chemical Toxicology*, 2011; 49(12):3025-30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22001171>

Mottier N, Jeanneret F, and Rotach M. Determination of hydrogen cyanide in cigarette mainstream smoke by lc/ms/ms. *Journal of AOAC International*, 2010; 93(3):1032–8. Available from: <http://www.atypon-link.com/AOAC/doi/pdf/10.5555/jaoi.93.3.1032?cookieSet=1>

International Agency for Research on Cancer. Monographs on the evaluation of carcinogenic risks to humans: Some naturally occurring substances: Food items and constituents, heterocyclic aromatic amines and mycotoxins., 56 Lyon, France: IARC, 1993. Available from: <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Some-Naturally-Occurring-Substances-Food-Items-And-Constituents-Heterocyclic-Aromatic-Amines-And-Mycotoxins-1993>.

International Agency for Research on Cancer. Agents classified by the IARC monographs, volumes 1–129. Lyon, France: IARC, Available from: <https://monographs.iarc.who.int/list-of-classifications>.

12.4.3.11 Metals, metalloids and radionuclides

Zhao, F, Chen, JX, Xu, H, Han, Y, Zhou, M, Wang, G et al. (2024). Arsenite Antiporter PvACR3 Driven by Its Native Promoter Increases Leaf Arsenic Accumulation in Tobacco. *Environ Sci Technol*, 58(26), 11534-11541. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38865317>

Ibrahim, AE, Alamir, SG, Al-Omairi, M, Salman, BI, Batakoushy, HA, Hegazy, MM, & Al-Harrasi, A. (2024). Tracking the Variations in Trace and Heavy Elements in Smoking Products Marketed in Oman and Egypt: Risk Assessment After Implementation of Constraining Protocols. *Biol Trace Elem Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38656680>

Su, X, Narayanan, M, Shi, X, Chen, X, Li, Z, & Ma, Y. (2024). Mitigating heavy metal accumulation in tobacco: Strategies, mechanisms, and global initiatives. *Sci Total Environ*, 926, 172128. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38565350>

Morillas, H, Gallego-Cartagena, E, & Upasen, S. (2024). Metals, nonmetals and metalloids in cigarette smoke as hazardous compounds for human health. *Sci Total Environ*, 923, 171351. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38432370>

Oner, F, Mazlumoglu, MR, Kurt, O, Kurt, N, & Ucuncu, H. (2024). Quantification of Heavy Metals in the Nasal Turbines of Smokers and Nonsmokers. *Laryngoscope*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38529719>

Alkufi, AA, Oleiwi, MH, & Abojassim, AA. (2024). Comparison of Heavy Metals in Urine Samples of Smoker and Non-smoker Persons. *Biol Trace Elem Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38386229>

Mashal, LA, Buss, FM, & Thabayneh, KM. (2024). (210)Po and (210)Pb radioactivity levels in local and imported tobacco used in Palestine and Jordan. *Radiat Environ Biophys*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38411924>

Farag, AGA, Badr, EA, Kholif, AOA, Khalifa, MN, & Ghanem, MMM. (2024). Serum and Seminal Plasma Levels of Lead and Arsenic in Cigarette Smokers and Their Relation to the Semen Parameters. *Biol Trace Elem Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38180596>

Felix, AT, & Ntarisa, AV. (2023). Review of natural radioactivity in tobacco cigarette brands. *J Environ Radioact*, 272, 107348. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38043217>

Hac, P, Rutkowska, M, Cieslik, BM & Konieczka, P. (2023). Estimation of smokers' exposure to mercury from combustible tobacco products, based on the approach used in food consumers' exposure estimation. *Food Chem Toxicol*, 114053. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37739052>

Hasan, M, Hossain, MM, Abrarin, S, Kormoker, T, Billah, MM, Bhuiyan, MKA et al. (2023). Heavy metals in popularly sold branded cigarettes in Bangladesh and associated health hazards from inhalation exposure. *Environ Sci Pollut Res Int*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37644270>

Piasek, M, Skrgatic, L, Sulimanec, A, Orct, T, Sekovanic, A, Kovacic, J et al. (2023). Effects of Maternal Cigarette Smoking on Trace Element Levels and Steroidogenesis in the Maternal-Placental-Fetal Unit. *Toxics*, 11(8) Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37624219>

Lin, W. (2023). Comparison of urine heavy metals in exclusive menthol and non-menthol cigarette users by race/ethnicity: The 2015-2016 National Health and Nutrition Examination Survey Special Sample. *Tob Prev Cessat*, 9, 22. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37389307>

Baghery, F, Lau, LDW, Mohamadi, M, Vazirinejad, R, Ahmadi, Z, Javedani, H et al. (2023). Risk of urinary tract cancers following arsenic exposure and tobacco smoking: a review. *Environ Geochem Health*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37248359>

Jia, XH, Li, YF, Fan, YG, Zhou, QH, Zhao, FH., Qiao, Y. L., & Lalley, M. (2023). Effect of Arsenic Exposure and Cigarette Smoking on Total and Cause-Specific Mortality: An Occupational Cohort With 27 Follow-up Years. *J Occup Environ Med*, 65(3), 217-223. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36868862>

Kim, B, Rhie, M, Park, S, Kim, HS, & Kwon, JA. (2023). Nonlinear Associations between Blood Cadmium Concentration and Thyroid Hormones According to Smoking Status in Korean Adults: The Korea National Health and Nutrition Examination Survey (KNHANES). *Toxics*, 11(2). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36851004>

Kim, J, Song, H, Lee, J, Kim, YJ, Chung, HS, Yu, JM et al. (2023). Smoking and passive smoking increases mortality through mediation effect of cadmium exposure in the United States. *Sci Rep*, 13(1), 3878. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36890267>

Jung, CC, Syu, ZH, Chou, CC, & Huang, YT. (2023). A study to characterize the lead isotopic fingerprint in PM(2.5) emitted from incense stick and cigarette burning. *Environ Sci Pollut Res Int*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36929257>

Sekovanic, A, Dorotic, A, Pasalic, D, Orct, T, Kljakovic-Gaspic, Z, Grgec, AS et al. (2022). The effects of maternal cigarette smoking on cadmium and lead levels, miRNA expression and biochemical parameters across the fetoplacental unit. *Heliyon*, 8(12), e12568. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36636214>

Day, PL, Wermers, M, Pazdernik, V, Jannetto, PJ, & Bornhorst, JA. (2022). Detection of Cadmium and Lead in Kidney Stones. Associations with Patient Demographics, Stone Composition, and Smoking. *J Appl Lab Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36575923>

Lau, LD, Willems, A, & Scardamaglia, L. (2022). Keratinocyte cancer in chronic smokers: is this arsenic exposure? *Med J Aust*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36458983>

Kozak, K, & Antosiewicz, DM. (2022). Tobacco as an efficient metal accumulator. *Biomaterials*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36097238>

Skalny AV, Serebryansky EP, Korobeinikova TV, Tsatsakis A, Vardavas C, et al. Smoking is associated with altered serum and hair essential metal and metalloid levels in women. *Food and Chemical Toxicology*, 2022:113249. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35728725>

Hac PJ, Cieslik BM, and Konieczka P. Review of cigars and cigar-type products as potential sources of consumer exposure to heavy metals. *J Environ Sci Health C Toxicol Carcinog*, 2022; 40(2):172-96. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35895917>

Al Mutairi MA, Al Herbish HA, Al-Ajmi RS, Alhazmi HZ, Al-Dhelaan RA, et al. Analyzing pesticides and metal(loid)s in imported tobacco to Saudi Arabia and risk assessment of inhalation exposure to certain metals. *Inhalation Toxicology*, 2022; 34(3-4):68-79. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35262446>

Vlachou C, Vejdovszky K, Wolf J, Steinwider J, Fuchs K, et al. Toxicological approaches for the quantitative inhalation risk assessment of toxic metals from tobacco smoke: Application on the deterministic and probabilistic inhalation risk assessment of cadmium for Austrian smokers. *Inhalation Toxicology*, 2021; 33(4):128-42. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33957849>

Salahel Din K. (210)pb and (210)po concentration levels in tobacco products and resulting radiation dose for Egyptian smokers. *Radiat Environ Biophys*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33730309>

Roselli C, Fagiolino I, Desideri D, Sisti D, and Meli MA. Assessment of the release of metals from cigarette butts into the environment. *PloS One*, 2021; 16(11):e0260111. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34793538>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8601425/pdf/pone.0260111.pdf>

Qamar W, Altamimi MA, Rehman MU, Ali N, Imam F, et al. Toxicological interaction between tobacco smoke toxicants cadmium and nicotine: An in-vitro investigation. *Saudi Journal of Biological Sciences*, 2021; 28(8):4201-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34354400>

Matassa R, Cattaruzza MS, Sandorfi F, Battaglione E, Relucenti M, et al. Direct imaging evidences of metal inorganic contaminants traced into cigarettes. *J Hazard Mater*, 2021; 411:125092. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33858086>

International Agency for Research on Cancer. IARC monographs on the evaluation of carcinogenic risks to humans: Arsenic, metals, fibres and dusts. 100C Lyon, France: IARC, 2021. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK304375/>.

Ho KJ, Chen TH, Yang CC, Chuang YC, and Chuang HY. Interaction of smoking and lead exposure among carriers of genetic variants associated with a higher level of oxidative stress indicators. *Int J Environ Res Public Health*, 2021; 18(16). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34444074>

Fireman Klein E, Klein I, Ephrat O, Dekel Y, Kessel A, et al. Trajectory of inhaled cadmium ultrafine particles in smokers. *BMJ Open Respir Res*, 2021; 8(1). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34845007>

Dinh QP, Novirsa R, Jeong H, Nugraha WC, Addai-Arhin S, et al. Mercury, cadmium, and lead in cigarettes from international markets: Concentrations, distributions and absorption ability of filters. *Journal of Toxicological Sciences*, 2021; 46(9):401-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34470992>

Van Duong H, Thanh Nguyen D, Peka A, Toth-Bodrogi E, and Kovacs T. 210po in soil and tobacco leaves in quang xuong, vietnam and estimation of annual effective dose to smokers. *Radiation Protection Dosimetry*, 2020; 192(1):106-12. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33230527>

Lisboa TP, Mimura AMS, da Silva JCJ, and de Sousa RA. Chromium levels in Tobacco, filter and ash of illicit brands cigarettes marketed in Brazil. *Journal of analytical toxicology*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31984423>

Lee JE, Kim HR, Lee MH, Kim NH, Wang KM, et al. Smoking-related DNA methylation is differentially associated with cadmium concentration in blood. *Biochem Genet*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32347401>

Ghanbar-Moghaddam B and Fathivand A. Study of polonium-210 in persian cigarette and Tobacco crops. *Radiat Prot Dosimetry*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33119076>

Fresquez MR, Watson CH, Valentin-Blasini L, and Pappas RS. Characterizing the transport of aluminum, silicon, and titanium-containing particles and nanoparticles in mainstream Tobacco smoke. *Journal of analytical toxicology*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33044491>

Ahmed AS, Aldubayan MA, Ahmed HA, Refaat AM, Alsalloumi AS, et al. Impact of smoking on heavy metal contamination and DNA fragmentation. *Environmental Science and Pollution Research International*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33201507>

Zhang P, Huang Z, Ma Y, Li Y, Ali N, et al. On-line detection of radioactive and non-radioactive heavy metals in tobacco smoke using portable laser-induced breakdown spectroscopy. *Analyst*, 2019; 144(11):3567-72. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31062786>

Repic A, Bulat P, Antonijevic B, Antunovic M, Dzudovic J, et al. The influence of smoking habits on cadmium and lead blood levels in the serbian adult people. *Environmental Science and Pollution Research International*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31811606>

Ozcan MM, Aljuhaimi F, Uslu N, Ghafoor K, Mohamed Ahmed IA, et al. Distribution of heavy metal and macroelements of Indian and imported cigarette brands in turkey. *Environmental Science and Pollution Research International*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31368072>

Kim K, Melough MM, Vance TM, Kim D, Noh H, et al. The relationship between zinc intake and cadmium burden is influenced by smoking status. *Food and Chemical Toxicology*, 2019; 125:210-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30615956>

Cheng LC, Lin CJ, Liu HJ, and Li LA. Health risk of metal exposure via inhalation of cigarette sidestream smoke particulate matter. *Environmental Science and Pollution Research International*, 2019; 26(11):10835-45. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30778946>

Ratelle M, Li X, and Laird BD. Cadmium exposure in first nations communities of the northwest territories, Canada: Smoking is a greater contributor than consumption of cadmium-accumulating organ meats. *Environ Sci Process Impacts*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30221302>

Ganguly K, Levanen B, Palmberg L, Akesson A, and Linden A. Cadmium in tobacco smokers: A neglected link to lung disease? *European Respiratory Review: An Official Journal of the European Respiratory Society*, 2018; 27(147). Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29592863>

Cuello-Nunez S, Benning J, Liu C, Branton P, Hu J, et al. Fractionation of cadmium in tobacco and cigarette smoke condensate using xanes and sequential leaching with icp-ms/ms. *Analytical and Bioanalytical Chemistry*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30094791>

Ates Alkan F, Karis D, Cakmak G, and Ercan AM. Analysis of the relationship between hemorheologic parameters, aluminum, manganese, and selenium in smokers. *Biol Trace Elem Res*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29704205>

Richter P, Faroon O, and Pappas RS. Cadmium and cadmium/zinc ratios and Tobacco-related morbidities. *Int J Environ Res Public Health*, 2017; 14(10). Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28961214>

Pinto E, Cruz M, Ramos P, Santos A, and Almeida A. Metals transfer from tobacco to cigarette smoke: Evidences in smokers' lung tissue. *J Hazard Mater*, 2017; 325:31-5. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27914289>

Liu H, Zhang Y, Zhou X, You X, Shi Y, et al. Source identification and spatial distribution of heavy metals in tobacco-growing soils in shandong province of China with multivariate and geostatistical analysis. *Environmental Science and Pollution Research International*, 2017; 24(6):5964-75. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28070814>

Benson NU, Anake WU, Adedapo AE, Fred-Ahmadu OH, and Ayejuyo OO. Toxic metals in cigarettes and human health risk assessment associated with inhalation exposure. *Environ Monit Assess*, 2017; 189(12):619. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29119337>

Aoki Y, Yee J, and Mortensen ME. Blood cadmium by race/hispanic origin: The role of smoking. *Environmental research*, 2017; 155:193-8. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28231546>

Tiwari M, Sahu SK, Bhangare RC, and Pandit GG. Polonium in size fractionated mainstream cigarette smoke, predicted deposition and associated internal radiation dose. *J Environ Radioact*, 2016; 162-163:251-7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27294663>

Pappas RS, Halstead MM, and Watson CH. Electron microscopic analysis of silicate and calcium particles in cigarette smoke tar. *Int J Respir Pulm Med*, 2016; 3(1). Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27158665>

Meltzer HM, Alexander J, Brantsaeter AL, Borch-Iohnsen B, Ellingsen DG, et al. The impact of iron status and smoking on blood divalent metal concentrations in Norwegian women in the hunt2 study. *J Trace Elem Med Biol*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27108098>

Kubalek D, Sersa G, Strok M, Benedik L, and Jeran Z. Radioactivity of cigarettes and the importance of (210)po and thorium isotopes for radiation dose assessment due to smoking. *J Environ Radioact*, 2016; 155-156:97-104. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26942842>

Rubio Armendariz C, Garcia T, Soler A, Gutierrez Fernandez AJ, Glez-Weller D, et al. Heavy metals in cigarettes for sale in Spain. *Environmental Research*, 2015; 143(Pt A):162-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26492401>

Pappas RS, Gray N, Gonzalez-Jimenez N, Fresquez M, and Watson CH. Triple quad-ICP-MS measurement of toxic metals in mainstream cigarette smoke from Spectrum Research cigarettes. *Journal of Analytical Toxicology*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26359486>

Afridi HI, Talpur FN, Kazi TG, and Brabazon D. Estimation of aluminum, arsenic, lead and nickel status in the samples of different cigarettes and their effect on human health of Irish smoker hypertensive consumers. *Clin Lab*, 2015; 61(9):1147-56. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26554233>

Matsunaga Y, Agaku IT, and Vardavas CI. The association between cigarette rod length, slim design, and blood cadmium levels among U.S. smokers: NHANES 1999–2010. *Prev Med*, 2014; 65:87-91. Available from: <http://www.sciencedirect.com/science/article/pii/S0091743514001522>

Caruso RV, O'Connor RJ, Stephens WE, Cummings KM, and Fong GT. Toxic metal concentrations in cigarettes obtained from U.S. smokers in 2009: Results from the International Tobacco Control (ITC) United States survey cohort. *Int J Environ Res Public Health*, 2014; 11(1):202-17. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24452255>

Arain SS, Kazi TG, Arain AJ, Afridi HI, Brahman KD, et al. Evaluated the levels of lead and cadmium in scalp hair of adolescent boys consuming different smokeless tobacco products with related to controls. *Biol Trace Elem Res*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25537077>

Arain SS, Gul Kazi T, Afridi HI, Brahman KD, NaEemuliah, et al. Arsenic content in smokeless tobacco products consumed by the population of Pakistan: Related health risk. *Journal of AOAC International*, 2014; 97(6):1662-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25632442>

Ajab H, Yaqub A, Malik SA, Junaid M, Yasmeen S, et al. Characterization of toxic metals in tobacco, tobacco smoke, and cigarette ash from selected imported and local brands in Pakistan. *ScientificWorldJournal*, 2014; 2014:413614. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24672317>

Afridi HI, Talpur FN, Kazi TG, and Brabazon D. Assessment of toxic elements in the samples of different cigarettes and their effect on the essential elemental status in the biological samples of Irish hypertensive consumers. *J Hum Hypertens*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25273861>

Takeuchi C, Correa-Afonso A, Pedrazzi H, Dinelli W, and Palma-Dibb R. Deposition of lead and cadmium released by cigarette smoke in dental structures and resin composite. *Microscopy Research and Technique*, 2010; [Epub ahead of print]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/jemt.20903/full>

Perez-Bernal J, Amigo J, Fernandez-Torres R, Bello M, and Callejon-Mochon M. Trace-metal distribution of cigarette ashes as marker of tobacco brands. *Forensic Science International*, 2010; 204(1-3):119-25. Available from:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20554412

Moeller D and Sun L. Chemical and radioactive carcinogens in cigarettes: Associated health impacts and responses of the tobacco industry, U.S. Congress, and federal regulatory agencies. *Health Physics*, 2010; 99(5):674–9. Available from: <http://journals.lww.com/health-physics/pages/articleviewer.aspx?year=2010&issue=11000&article=00009&type=abstract>

Afridi H, Kazi T, Kazi N, Jamali M, Arain M, et al. Evaluation of cadmium, lead, nickel and zinc status in biological samples of smokers and nonsmokers hypertensive patients. *Journal of Human Hypertension*, 2010; 24(1):34–43. Available from: <http://www.nature.com/jhh/journal/v24/n1/full/jhh200939a.html>

Papastefanou C. Radioactivity of tobacco leaves and radiation dose induced from smoking. *Int J Environ Res Public Health*, 2009; 6(2):558–67. Available from: <http://www.mdpi.com/1660-4601/6/2/558>

Massadeh A, Gharibeh A, Omari K, Al-Momani I, Alomari A, et al. Simultaneous determination of cd, pb, cu, zn, and se in human blood of jordanian smokers by icp-oes. *Biological Trace Element Research*, 2009; 133(1):1–11. Available from: <http://www.springerlink.com/content/7020123023174710/fulltext.html>

Martin C, Antonini J, and Doney B. A case report of elevated blood cadmium. *Occupational Medicine*, 2009; 59(2):130–2. Available from: <http://occmmed.oxfordjournals.org/cgi/content/full/59/2/130>

Kazi T, Wadhwa S, Afridi H, Kazi N, Kandhro G, et al. Interaction of cadmium and zinc in biological samples of smokers and chewing tobacco female mouth cancer patients. *Journal of Hazardous Materials*, 2009; 176(1–3):985–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20031315>

Cox L. Could removing arsenic from tobacco smoke significantly reduce smoker risks of lung cancer? *Risk Analysis* 2008; 29(1):3–17. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1539-6924.2008.01145.x/full>

Tso TC, Harley N, and Alexander LT. Source of lead-210 and polonium-210 in tobacco. *Science*, 1966; 153(3738):880-2. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/5914751>

12.4.3.12 Additives and flavours (refers to section 12.6)

12.4.4 Emission from tobacco that is not combusted (set alight)

Cao, Y, Liu, X, Hu, Z, Li, J, Chen, X, Xiong, Y et al. (2024). Assessing nicotine pharmacokinetics of new generation tobacco products and conventional cigarettes: a systematic review and meta-analysis. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39162577>

12.4.4.1 Emissions from waterpipes

Kassem, NOF, Strongin, RM, Stroup, AM, Brinkman, MC, El-Hellani, A, Erythropel, HC et al. (2024). Toxicity of Waterpipe Tobacco Smoking: The Role of Flavors, Sweeteners, Humectants, and Charcoal. *Toxicol Sci*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39037923>

Almomen, S, Aldossari, M, Khaleel, Y, Altamimi, M, Alharbi, O, Alsuwaydani, A et al. (2023). Effect of glycerol concentration on levels of toxicants emissions from water-pipe tobacco smoking (WTS). *BMC Public Health*, 23(1), 1858. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37749520>

Abadi, DRV, Tahmasbizadeh, M, Arfaeina, H, Masjedi, MR, Ramavandi, B, & Poureshgh, Y. (2022). Biomonitoring of unmetabolized polycyclic aromatic hydrocarbons (PAHs) in urine of waterpipe/cigarette cafe workers. *Environ Sci Pollut Res Int*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36306072>

Bakker-‘t Hart IME, Bakker F, Pennings JLA, Weibolt N, Eising S, et al. Flavours and flavourings in waterpipe products: A comparison between tobacco, herbal molasses and steam stones. *Tobacco Control*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35241500>

Jaccard G, Tabin Djoko D, Korneliou A, and Belushkin M. Analysis of waterpipe aerosol constituents in accordance with the ISO standard 22486. *Toxicology Reports*, 2020; 7:1344-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33102137>

El Hourani M, Salman R, Talih S, Saliba NA, and Shihadeh A. Does the bubbler scrub key toxicants from waterpipe tobacco smoke?: Measurements and modeling of co, no, pah, nicotine, and particulate matter uptake. *Chemical Research in Toxicology*, 2020; 33(3):727-30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31957423>

Rezk-Hanna M and Benowitz NL. Cardiovascular effects of hookah smoking: Potential implications for cardiovascular risk. *Nicotine & Tobacco Research*, 2019; 21(9):1151-61. Available from: <https://pubmed.ncbi.nlm.nih.gov/29660041/>

El Hourani M, Talih S, Salman R, Karaoghlanian N, Karam E, et al. Comparison of co, pah, nicotine, and aldehyde emissions in waterpipe tobacco smoke generated using electrical and charcoal heating methods. *Chemical Research in Toxicology*, 2019; 32(6):1235-40. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31038931>

Shihadeh A, Schubert J, Klaiany J, El Sabban M, Luch A, et al. Toxicant content, physical properties and biological activity of waterpipe tobacco smoke and its tobacco-free alternatives. *Tobacco Control*, 2015; 24 Suppl 1:i22-i30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25666550>

Schubert J, Hahn J, Dettbarn G, Seidel A, Luch A, et al. Mainstream smoke of the waterpipe: Does this environmental matrix reveal as significant source of toxic compounds? *Toxicology letters*, 2011; 205(3):279-84. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21712083>

Cobb C, Ward KD, Maziak W, Shihadeh AL, and Eissenberg T. Waterpipe tobacco smoking: An emerging health crisis in the United States. *American Journal of Health Behavior*, 2010; 34(3):275-85. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20001185>

Monzer B, Sepetdjian E, Saliba N, and Shihadeh A. Charcoal emissions as a source of co and carcinogenic pah in mainstream narghile waterpipe smoke. *Food and Chemical Toxicology*, 2008; 46(9):2991-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18573302>

Shihadeh A. Investigation of mainstream smoke aerosol of the argileh water pipe. *Food and Chemical Toxicology*, 2003; 41(1):143-52. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12453738>

12.4.4.2 Emissions from heated tobacco ('heat-not-burn') products

Wen, Z, Li, X, Gu, X, Zhang, W, Pang, Y, Jiang, X et al. (2022). Online analysis of chemical composition and size distribution of fresh cigarette smoke emitted from a heated tobacco product. *MethodsX*, 9, 101912. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36385917>

Salman R, Talih S, El-Hage R, Haddad C, Karaoghlanian N, et al. Free-base and total nicotine, reactive oxygen species, and carbonyl emissions from iqos, a heated tobacco product. *Nicotine & Tobacco Research*, 2019; 21(9):1285-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30476301>

Davis B, Williams M, and Talbot P. Iqos: Evidence of pyrolysis and release of a toxicant from plastic. *Tobacco Control*, 2019; 28(1):34-41. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29535257>

Leigh NJ, Palumbo MN, Marino AM, O'Connor RJ, and Goniewicz ML. Tobacco-specific nitrosamines (tsna) in heated tobacco product iqos. *Tobacco Control*, 2018; 27(Suppl 1):s37-s8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30242043>

Farsalinos KE, Yannovits N, Sarri T, Voudris V, and Poulas K. Nicotine delivery to the aerosol of a heat-not-burn tobacco product: Comparison with a tobacco cigarette and e-cigarettes. *Nicotine & Tobacco Research*, 2018; 20(8):1004-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28637344>

Eaton D, Jakaj B, Forster M, Nicol J, Mavropoulou E, et al. Assessment of tobacco heating product thp1.0. Part 2: Product design, operation and thermophysical characterisation. *Regulatory Toxicology and Pharmacology*, 2018; 93:4-13. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29080851>

Food & Drug Administration. Pmta cover sheet: Technical project lead review for iqos Tobacco heating system (ths). FDA, 2017. Available from: <https://www.fda.gov/media/124247/download>.

Auer R, Concha-Lozano N, Jacot-Sadowski I, Cornuz J, and Berthet A. Heat-not-burn tobacco cigarettes: Smoke by any other name. *JAMA Internal Medicine*, 2017; 177(7):1050-2. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28531246>

Lopez AA, Hiler M, Maloney S, Eisenberg T, and Breland AB. Expanding clinical laboratory tobacco product evaluation methods to loose-leaf tobacco vaporizers. *Drug and Alcohol Dependence*, 2016; 169:33-40. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27768968>

Forster M, Liu C, Duke MG, McAdam KG, and Proctor CJ. An experimental method to study emissions from heated tobacco between 100-200 degrees c. *Chemistry Central Journal*, 2015; 9:20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25941536>

Stapleton JA, Russell MAH, Sutherland G, and Feyerabend C. Nicotine availability from eclipse tobacco-heating cigarette. *Psychopharmacology*, 1998; 139(3):288–90. Available from: <http://www.springerlink.com/content/xpj7bt8hbl3m3c8r/>

12.4.4.3 Emission from smokeless tobacco

Edwards, SH, Hassink, M, Taylor, KM, & Vu, AT. (2023). Variation of Benzo[a]pyrene, NNN, and NNK Levels in 16 Commercial Smokeless Tobacco Products. *Chem Res Toxicol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36638203>

Wilhelm J, Mishina E, Viray L, Paredes A, and Pickworth WB. The pH of smokeless tobacco determines nicotine buccal absorption: Results of a randomized crossover trial. *Clinical Pharmacology and Therapeutics*, 2022; 111(5):1066-74. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34826137>

Xia B, Blount BC, Guillot T, Brosius C, Li Y, et al. Tobacco-specific nitrosamines (NNAL, nnn, nat, and nab) exposures in the US population assessment of Tobacco and health (path) study wave 1 (2013-2014). *Nicotine & Tobacco Research*, 2021; 23(3):573-83. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32716026>

Miller JH, Danielson T, Pithawalla YB, Brown AP, Wilkinson C, et al. Method development and validation of dissolution testing for nicotine release from smokeless tobacco products using flow-through cell apparatus and uplc-pda. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences*, 2020; 1141:122012. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32065955>

Cheng YC, Reyes-Guzman CM, Christensen CH, Rostron BL, Edwards KC, et al. Biomarkers of exposure among adult smokeless tobacco users in the population assessment of Tobacco and health study (wave 1, 2013-2014). *Cancer Epidemiology, Biomarkers & Prevention*, 2020; 29(3):659-67. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31988072>

Prasad GL, Jones BA, Chen P, and Gregg EO. A cross-sectional study of biomarkers of exposure and effect in smokers and moist snuff consumers. *Clinical Chemistry and Laboratory Medicine*, 2016; 54(4):633-42. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26495926>

Hatsukami DK, Stepanov I, Severson H, Jensen JA, Lindgren BR, et al. Evidence supporting product standards for carcinogens in smokeless tobacco products. *Cancer Prevention Research (Philadelphia, Pa.)*, 2015; 8(1):20-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25524878>

McAdam KG, Faizi, A., Kimpton, H., Porter, A., Rodu, B. Polycyclic aromatic hydrocarbons in US and Swedish smokeless tobacco products. *Chemistry Central Journal*, 2013; 7.

Stepanov I, Villalta P, Knezevich A, Jensen J, Hatsukami D, et al. Analysis of 23 polycyclic aromatic hydrocarbons in smokeless tobacco by gas chromatography-mass spectrometry. *Chemical Research in Toxicology*, 2010; 23(1):66-73. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19860436>

Boffetta P and Straif K. Use of smokeless tobacco and risk of myocardial infarction and stroke: Systematic review with meta-analysis. *British Medical Journal (Clinical Research ed)*, 2009; 339:b3060. Available from: <http://www.ncbi.nlm.nih.gov/entrez/pubmed19690343>

Stepanov I, Jensen J, Hatsukami D, and Hecht S. New and traditional smokeless tobacco: Comparison of toxicant and carcinogen levels. *Nicotine & Tobacco Research*, 2008;

10(12):1773-82. Available from:

<http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200802443544>

Stepanov I, Jensen J, Hatsukami D, and Hecht SS. Tobacco-specific nitrosamines in new tobacco products. *Nicotine & Tobacco Research*, 2006; 8(2):309-13. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/16766423>

McNeill A, Bedi R, Islam S, Alkhatib MN, and West R. Levels of toxins in oral tobacco products in the UK. *Tobacco Control*, 2006; 15(1):64-7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/16436408>

World Health Organization and International Agency for Research on Cancer. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 85: Betel-quid and areca-nut chewing and some areca-nut-derived nitrosamines. Summary of data reported and evaluation. Lyon: IARC, 2004. Available from:

<http://monographs.iarc.fr/ENG/Monographs/vol85/volume85.pdf>.

Chapman S and Wakefield M. Tobacco control advocacy in Australia: Reflections on 30 years of progress. *Health Education and Behavior*, 2001; 28(3):274-89. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/11380049>

Tomar SL and Henningfield JE. Review of the evidence that pH is a determinant of nicotine dosage from oral use of smokeless tobacco. *Tobacco Control*, 1997; 6(3):219-25. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/9396107>

Brunnemann KD and Hoffmann D. Chemical composition of smokeless tobacco products in Smokeless tobacco or health: An international perspective. Smoking and tobacco control monograph no. 2 Bethesda, MA: National Cancer Institute, Department of Health and Human Services; 1992. Available from: https://cancercontrol.cancer.gov/sites/default/files/2020-08/m02_complete.pdf.

Benowitz NL. Pharmacology of smokeless tobacco use: Nicotine addiction and nicotine-related health consequences, in Smokeless tobacco or health: An international perspective. Smoking and tobacco control monograph no. 2 Bethesda, MA: National Cancer Institute, Department of Health and Human Services; 1992. Available from: https://cancercontrol.cancer.gov/sites/default/files/2020-08/m02_complete.pdf.

News:

12.4 Emissions from tobacco products

World Health Organization. Report on the scientific basis of tobacco product regulation: Fifth report of a WHO study group. WHO, 2015. Last update: Viewed Available from:

<https://apps.who.int/iris/bitstream/handle/10665/161512/9789241209892.pdf?sequence=1&isAllowed=y>.

World Health Organization. Partial guidelines for implementation of articles 9 and 10 of the WHO Framework Convention on Tobacco Control. 2013. Last update: Viewed Available from:

https://www.who.int/fctc/guidelines/Guidelines_Articles_9_10_rev_240613.pdf.

Administration FaD. Harmful and potentially harmful constituents in Tobacco products and Tobacco smoke: Established list Federal Register / Vol. 77, No. 64 / Tuesday, April 3, 2012 / Notices Silver Spring, Maryland: Food and Drug Administration, U.S. Department of Health and Human Services, 2012. Last update: 07/07/2017 Viewed 25/07/2017. Available from:

<http://www.fda.gov/downloads/TobaccoProducts/Labeling/RulesRegulationsGuidance/UCM297981.pdf>.

12.4.1 What is tobacco smoke?

Mohammadi D. Guilty pleasures: Just how bad is social smoking?, in New Scientist 2015: London: UK. Available from: <https://www.newscientist.com/article/mg22630233-800-guilty-pleasures-just-how-bad-is-social-smoking/>.

Ontario Ministry of Health, Collaborative Change, and BBDO. Ontario ministry of health: Social smoking. Collaborative Change 2013. Last update: Unknown; Viewed 14 September 2018. Available from: http://www.collaborativechange.org.uk/collaborative-change-tackle-social-smoking/?doing_wp_cron=1536449645.9569699764251708984375

12.4.1.1 Particles and gases in tobacco smoke

12.4.1.2 Mainstream smoke, sidestream smoke and secondhand smoke

Mohammadi D. Guilty pleasures: Just how bad is social smoking?, in New Scientist 2015: London: UK. Available from: <https://www.newscientist.com/article/mg22630233-800-guilty-pleasures-just-how-bad-is-social-smoking/>.

Ontario Ministry of Health, Collaborative Change, and BBDO. Ontario ministry of health: Social smoking. Collaborative Change 2013. Last update: Unknown; Viewed 14 September 2018. Available from: http://www.collaborativechange.org.uk/collaborative-change-tackle-social-smoking/?doing_wp_cron=1536449645.9569699764251708984375.

12.4.2 What happens when tobacco products are burned?

12.4.2.1 Pyrolysis

12.4.2.2 Combustion

12.4.2.3 Formation of smoke in cigarettes

12.4.2.4 Formation of smoke in other types of combusted tobacco products

Mottola G. 10 things every cigar smoker should know. 2017. Last update: Viewed Available from: <https://www.cigaraficionado.com/article/10-things-every-cigar-smoker-should-know-19531>.

12.4.3 Chemicals that are emitted from combusted tobacco products

Bevan L. 7 poisons you consume every time you smoke, and you thought it's just Tobacco. The Epoch Times, 2019. Available from: https://www.theepochtimes.com/7-poisons-you-consume-every-time-you-smoke-and-you-thought-its-just-tobacco_2975549.html

Food and Drug Administration. Harmful and potentially harmful constituents in tobacco products and tobacco smoke: Established list. 2012. Last update: Viewed Available from: <https://www.fda.gov/tobacco-products/rules-regulations-and-guidance/harmful-and-potentially-harmful-constituents-tobacco-products-and-tobacco-smoke-established-list>.

Federal Trade Commission. Tar, nicotine and carbon monoxide of the smoke of 1252 varieties of domestic cigarette for the year 1997. 1999. Last update: Viewed accessed September 1999. Available from: www.ftc.gov.

National Cancer Institute. Tobacco tar. NCI Dictionaries: NCI, Last update: Viewed Available from: <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/tobacco-tar>.

12.4.3.1 Nicotine and other alkaloids

Rogers K. Alkaloid. Britannica, 2002. Last update: Viewed Available from: <https://www.britannica.com/science/alkaloid>.

12.4.3.2 Aldehydes

Stainton LH. Scientists help id new cancer-causing agent in Tobacco smoke. NJ Spotlight, 2018. Available from: <http://www.njspotlight.com/stories/18/06/21/rutgers-scientist-helps-id-new-cancer-causing-agent-in-tobacco-smoke/>

Food and Drug Administration. Harmful and potentially harmful constituents in tobacco products and tobacco smoke: Established list. 2012. Last update: Viewed Available from: <https://www.fda.gov/tobacco-products/rules-regulations-and-guidance/harmful-and-potentially-harmful-constituents-tobacco-products-and-tobacco-smoke-established-list>.

Rogers K. Aldehyde: Chemical compound. Britannica, 2008. Last update: Viewed Available from: <https://www.britannica.com/science/aldehyde>.

The Editors of Encyclopaedia Britannica. Formaldehyde: Chemical compound. Britannica, 1998. Last update: Viewed Available from: <https://www.britannica.com/science/formaldehyde>.

12.4.3.3 Aromatic amines

12.4.3.4 Hydrocarbons

Carey FA and Rogers K. Hydrocarbon: Chemical compound. Britannica, 2008. Last update: Viewed Available from: <https://www.britannica.com/science/hydrocarbon>.

12.4.3.5 Heterocyclic aromatic amines

Heterocyclic amine. Comprehensive toxicology 2010. Last update: Viewed Available from: <https://www.sciencedirect.com/topics/medicine-and-dentistry/heterocyclic-amine>.

12.4.3.6 Polycyclic aromatic hydrocarbons (PAH)

Centers for Disease Control and Prevention. Polycyclic aromatic hydrocarbons (pahs) factsheet. 2017. Last update: Viewed Available from: https://www.cdc.gov/biomonitoring/PAHs_FactSheet.html.

12.4.3.7 N-nitrosamines

Centers for Disease Control and Prevention. NNAL (4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol) factsheet. CDC, 2017. Last update: Viewed Available from: https://www.cdc.gov/biomonitoring/NNAL_FactSheet.html.

12.4.3.8 Phenols

12.4.3.9 Other carbon-based (organic) compounds

Food & Drug Administration. Food for human consumption (continued) part 189 -- substances prohibited from use in human food. 2020. Last update: Viewed Available from: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcr/CFRSearch.cfm?fr=189.130>.

Centers for Disease Control and Prevention. Acrylamide factsheet. CDC, 2017. Last update: Viewed Available from: https://www.cdc.gov/biomonitoring/Acrylamide_FactSheet.html.

12.4.3.10 Inorganics

12.4.3.11 Metals, metalloids and radionuclides

12.4.3.12 Additives and flavours (refers to section 12.6)

12.4.4 Emission from tobacco that is not combusted (set alight)

12.4.4.1 Emissions from waterpipes

12.4.4.2 Emissions from heated tobacco ('heat-not-burn') products

World Health Organization. Heated tobacco products information sheet. WHO, 2020. Viewed:
Available from: <https://www.who.int/publications/i/item/WHO-HEP-HPR-2020.2>.

Philip Morris International. Iqos heated tobacco products. Last update: Viewed Available from:
<https://www.pmi.com/smoke-free-products/iqos-our-tobacco-heating-system>.

Ploom UK. What are heated tobacco / t-vapour products? : JTI UK, Last update: Viewed Available
from: <https://www.ploom.co.uk/faq/9/>.

12.4.4.3 Emission from smokeless tobacco

Australian Competition & Consumer Commission. Smokeless tobacco products. Last update: Viewed
Available from: <https://www.productsafety.gov.au/products/health-lifestyle/personal/tobacco-related-products/smokeless-tobacco-products>.

World Health Organization. Report on the scientific basis of tobacco product regulation: Fifth report
of a WHO study group. WHO, 2015. Last update: Viewed Available from:
<https://apps.who.int/iris/bitstream/handle/10665/161512/9789241209892.pdf?sequence=1&isAllowed=y>.